arXiv:1112.0397v2 [math.DS] 11 Dec 2011

A Geometric Path from Zero Lyapunov Exponents to Rotation Cocycles

Jairo Bochi* & Andrés Navas†

December 2, 2011

Abstract

We consider cocycles of isometries on spaces of nonpositive curvature H. We show that if such a cocycle has uniform sublinear drift, then there are almost invariant sections, i.e., continuous sections that move very little under the cocycle dynamics. The construction is based on an appropriate concept of barycenter in H. If, in addition, H is a symmetric space, then we show that almost invariant sections can be made invariant by perturbing the cocycle. Applying these results to the symmetric space $H = \mathrm{GL}(d, \mathbb{R})/\mathrm{O}(d)$, we conclude that if all Lyapunov exponents of a matrix cocycle vanish, then it can be approximated by a cocycle that is cohomologous to a cocycle taking values in the orthogonal group $\mathrm{O}(d)$. This extends a result of Avila, Bochi and Damanik to more general dynamics and arbitrary cocycle dimension. For linear cocycles with non-necessarily vanishing Lyapunov exponents, we show that the property of existence of conformal invariant structures along the Oseledets splitting is dense in the space of continuous cocycles.

---

*Partially supported by CNPq (Brazil) and the Math-AMSUD Project DySET.

†Partially supported by CONICYT (Chile) and the Math-AMSUD Project DySET.

Table of contents

1 Introduction 3
- 1.1 From subexponential growth to boundedness . . . 3
- 1.2 Sketch of the proof of Theorem 1.1 . . . 4
- 1.3 The general main results: discrete time . . . 6
- 1.4 The general main results: continuous time . . . 8
- 1.5 Invariant conformal structures for matrix cocycles . . . 10
- 1.6 Further remarks and questions . . . 10
- 1.7 Organization of the paper . . . 12

2 Proof of the results for discrete-time cocycles 12
- 2.1 Busemann spaces . . . 12
- 2.2 Barycenter maps . . . 13
- 2.3 Proof of Theorem A . . . 14
- 2.4 Isometries . . . 15
- 2.5 Symmetric spaces and transvections . . . 15
- 2.6 Curvature bounds . . . 16
- 2.7 Macroscopic uniform homogeneity . . . 17
- 2.8 Proof of Theorem B . . . 18
- 2.9 Proof of Theorem 1.1 . . . 19

3 Proof of the results for continuous-time cocycles 20
- 3.1 Cartan–Hadamard manifolds . . . 20
- 3.2 Killing fields and symmetric manifolds . . . 20
- 3.3 Infinitesimal displacement estimates . . . 21
- 3.4 Infinitesimal uniform homogeneity . . . 22
- 3.5 Proof of Theorem C . . . 23
- 3.6 Proof of Theorem D . . . 24

4 Fiber bundle versions 25

5 Invariant conformal structures 27

A Appendix: The displacement estimate for non-proper spaces 29
- A.1 Angles and more comparisons . . . 29
- A.2 The displacement estimate . . . 30

B Appendix: Some lemmas on Killing fields 32

C Appendix: Differentiability of the Cartan barycenter 33

References 34

1 Introduction

1.1 From subexponential growth to boundedness

A *cocycle* over a (continuous) map $F\colon \Omega \to \Omega$ is a continuous function

$$A\colon \Omega \to \mathbb{G}, \tag{1}$$

where Ω is a compact Hausdorff topological space and $\mathbb{G}$ is a topological group. We denote $A^{(0)}(\omega) := \mathrm{id}$ and for $n \in \mathbb{Z}_+$,

$$A^{(n)}(\omega) := A(F^{n-1}\omega) \cdots A(F\omega)A(\omega). \tag{2}$$

Notice the *cocycle relation*

$$A^{(n+m)}(\omega) = A^{(m)}(F^n\omega)A^{(n)}(\omega). \tag{3}$$

Two cocycles A and B (over F) are said to be *cohomologous* whenever there exists a continuous map $U\colon \Omega \to \mathbb{G}$ such that

$$A(\omega) = U(F\omega)B(\omega)U(\omega)^{-1}, \quad \text{for all } \omega \in \Omega. \tag{4}$$

Now suppose that $\mathbb{G}$ is the group $\mathrm{GL}(d,\mathbb{R})$ of invertible $d \times d$ real matrices. Then a cocycle $A\colon \Omega \to \mathrm{GL}(d,\mathbb{R})$ is called a *matrix cocycle* (or a *linear cocycle*).

We say that the linear cocycle A has *uniform subexponential growth* if

$$\lim_{n\to+\infty} \frac{1}{n} \log \left\| [A^{(n)}(\omega)]^{\pm 1} \right\| = 0 \quad \text{uniformly over } \omega \in \Omega, \tag{5}$$

for some (and hence any) matrix norm $\|\cdot\|$. (In fact, this is equivalent to the vanishing of all Lyapunov exponents with respect to all F-invariant probability measures: see Remark 1.3.)

We say that A is *product-bounded*[1] if

$$\sup_{\omega\in\Omega,\ n\in\mathbb{Z}_+} \left\| [A^{(n)}(\omega)]^{\pm 1} \right\| < \infty \tag{6}$$

If a cocycle A is cohomologous to a cocycle B taking values in the orthogonal group $\mathrm{O}(d)$, then A is product-bounded, as it is easy to check. Although product-bounded cocycles are not always cohomologous to cocycles with values in $\mathrm{O}(d)$ (see e.g. [KatH, Exercise 2.9.2], [Qu]), this happens whenever F is minimal, according to a result shown by Coronel, Navas and Ponce in [CNP1]. (In the non-minimal case, one can still ensure the existence of a bounded and measurable conjugacy U.)

Our first result reads as follows:

---

[1] One should avoid confusion with the other concept of product-boundedness (studied e.g. in [Mo]) where one does not take inverses in (6).

Theorem 1.1. *Let A be a matrix cocycle with uniform subexponential growth. Then there exists $\tilde{A}\colon \Omega \to \mathrm{GL}(d, \mathbb{R})$ arbitrarily close to A such that $\tilde{A}$ is cohomologous to a cocycle taking values in $\mathrm{O}(d)$.*

Here, closeness is meant with respect to the uniform (C^0) topology on the space of continuous maps from Ω into $\mathrm{GL}(d, \mathbb{R})$.

For maps A taking values in the group $\mathrm{SL}(2, \mathbb{R})$ and under extra assumptions on the dynamics F, the result above was shown by Avila, Bochi and Damanik as a step in the proofs of their results about spectra of Schrödinger operators, see [ABD1, ABD2].[2]

As we will see, our method of proof also allows us to obtain versions of Theorem 1.1 for other matrix groups, as the complex general linear group and the symplectic group, and even certain groups of linear operators in infinite dimension; see Remarks 2.12 and 2.13.

The rest of this Introduction goes as follows. In § 1.2, we explain the ideas behind the proof of Theorem 1.1. These ideas rely on the fact that invertible matrices act by isometries on a natural metric space H; then the proof consists on purely geometrical constructions on H. In §§ 1.3 and 1.4, we state our main results, which concern cocycles of isometries of much more general spaces H (and actually imply Theorem 1.1). In § 1.5, we come back to general matrix cocycles. Combining the techniques involved in the proof of Theorem 1.1 with a result from Bochi and Viana [BV], we show that every matrix cocycle can be perturbed so that its action on the bundles of the finest dominated splitting become conformal.

1.2 Sketch of the proof of Theorem 1.1 and hints of generalization

Consider cocycles in the group $\mathbb{R}$. In this case, we will use the letter g instead of A, and in additive notation the product from (2) becomes a Birkhoff sum:

$$g^{(n)}(\omega) := g(\omega) + \cdots + g(F^{n-1}\omega), \qquad g^{(0)}(\omega) := 0.$$

So two continuous functions g, $h\colon \Omega \to \mathbb{R}$ are cohomologous (with respect to the dynamics F) if there is a continuous function $\varphi\colon \Omega \to \mathbb{R}$ such that

$$g = \varphi \circ F + h - \varphi.$$

If h is identically zero, then g is called a *coboundary.*

Using the fact that $\mathrm{GL}_+(1, \mathbb{R}) \sim \mathbb{R}$, the case $d = 1$ in Theorem 1.1 reduces to the following result:

Theorem 1.2 (Baby Theorem: The one-dimensional case)**.** *Let $g\colon \Omega \to \mathbb{R}$ be a continuous function. If the Birkhoff averages $\frac{1}{n} g^{(n)}$ converge uniformly to zero, then there exists a coboundary $\tilde{g}$ arbitrarily close to g.*

---

[2] However, they haven't explicitly stated the result: see the proof of Theorem 1 in [ABD1] and Proposition 6.3 in [ABD2].

Proof. [3] Define a sequence of continuous functions $\varphi_N \colon \Omega \to \mathbb{R}$ by

$$\varphi_N(\omega) = \frac{1}{N} \sum_{i=0}^{N-1} [-g^{(i)}(\omega)]. \tag{7}$$

Then

$$\varphi_N(F\omega) = \frac{1}{N} \sum_{i=0}^{N-1} [-g^{(i)}(F\omega)] = \frac{1}{N} \sum_{i=0}^{N-1} \left[-g^{(i+1)}(\omega) + g(\omega) \right]$$
$$= \varphi_N(\omega) + g(\omega) + \frac{g^{(N)}(\omega)}{N}\,.$$

Using the hypothesis, we conclude that the sequence of coboundaries $\varphi_N \circ F - \varphi_N$ converges uniformly to g. ☐

We reread the proof above as follows: Consider the skew-product map

$$G = G_{F,g} \colon \Omega \times \mathbb{R} \to \Omega \times \mathbb{R} \quad \text{given by} \quad (\omega, s) \mapsto (F(\omega), s + g(\omega)). \tag{8}$$

Then for large N, the function $\varphi = \varphi_N \colon \Omega \to \mathbb{R}$ is an *almost invariant section*, in the sense that the G-image of its graph is very close to itself. What we did was to modify g to $\tilde{g}$ so that the associated skew-product $G_{F,\tilde{g}}$ leaves the graph of φ exactly invariant. This in turn implies that $\tilde{g}$ is cohomologous to the trivial cocycle, i.e., is a coboundary.

Now assume $d \geq 2$. Let $\mathcal{E} = \mathcal{E}_d$ be the space of ellipsoids in $\mathbb{R}^d$ centered at the origin. The group $\mathrm{GL}(d, \mathbb{R})$ acts on $\mathcal{E}$ in the obvious way, and this action is by isometries for a certain natural distance on the space $\mathcal{E}$. Moreover, for this distance d, a cocycle A has uniform subexponential growth if and only if

$$\lim_{n \to \infty} \frac{1}{n} d\big(A^{(n)}(\omega)e, e\big) = 0 \quad \text{uniformly over } \omega \in \Omega \tag{9}$$

for some (and hence any) $e \in \mathcal{E}$. (See § 2.9.)

Given a linear cocycle A, consider the skew-product map $G = G_{F,A} \colon \Omega \times \mathcal{E} \to \Omega \times \mathcal{E}$ given by $G(\omega, e) = (F(\omega), A(\omega)e)$. Suppose that the map T has an invariant section, that is, a continuous map $\varphi \colon \Omega \to \mathcal{E}$ whose graph is T-invariant. It is possible to find a continuous map $U \colon \Omega \to \mathrm{GL}(d, \mathbb{R})$ such that $\varphi(\omega) = U(\omega)^{-1} e_0$, where e_0 is the unit ball in $\mathbb{R}^d$. Let $B(\omega) = U(F\omega)A(\omega)U(\omega)^{-1}$. The invariance property $A(\omega)\varphi(\omega) = \varphi(F\omega)$ translates to $B(\omega)e_0 = e_0$. Thus B takes values in $\mathrm{O}(d)$.

Thus to prove Theorem 1.1, it suffices to perturb A to $\tilde{A}$ so that the associated skew-product has an invariant section. We can do this in two steps:

---

[3] This argument seems to be folklore. See [MOP2]; see also the proof of Proposition 6 of [CNP2]. For an indirect proof of Theorem 1.2 (using the Hahn–Banach theorem), see [Kat, Prop. 2.13].

1. We first show that if (F, A) has uniform subexponential growth, then there is a sequence of almost invariant sections $\varphi_N \colon \Omega \to \mathcal{E}$; more precisely, the distance between the graph of φ_N and its F-image tends to zero as $N \to \infty$.

2. Next we show that if (F, A) has a almost invariant section φ, then one can perturb A so that φ becomes invariant.

Let's give clues on how to perform each of the steps:

1. For Step 1, the idea to construct almost invariant sections φ_N is to imitate formula (7), replacing the averaging operation $\frac{1}{N}\sum_{i=0}^{N-1}$ by an appropriate *barycenter* concept.

2. In Step 2, the perturbation $\tilde{A}(\omega)$ of $A(\omega)$ is obtained by left-multiplying $A(\omega)$ by a matrix close to the identity that takes the ellipsoid $A(\omega)\varphi(\omega)$ to the nearby ellipsoid $\varphi(F\omega)$. This is not as trivial as it seems, because these two ellipsoids may be very distorted, and moreover the correcting matrices must be chosen in a way that depends continuously on ω. We accomplish this by establishing a certain *uniform homogeneity* property of the space of ellipsoids.

These two properties, existence of a good barycenter concept and uniform homogeneity, are purely geometrical. In fact, not only the space of ellipsoids but much more general spaces posses these properties. Besides simple connectedness, the essential ingredient for both properties is *nonpositive curvature.* (Also, for the homogeneity property, it is necessary that the space has enough isometries.) Theorem 1.1 will then follow from a much more general result for cocycles taking values in the group of isometries of a nonpositively curved space. We remark that this approach comes from the seminal work of Karlsson and Margulis [KarM], who proved a geometric version of the Oseledets' theorem in this setting. (See also [Kai, KarL].)

1.3 The general main results: discrete time

Let H be a Busemann space (i.e., a separable complete geodesic space of nonpositive curvature in the sense of Busemann), and let $\mathbb{G} := \mathrm{Isom}(H)$ be its group of isometries, endowed with the bounded–open topology. (Definitions are given in § 2.1.)

We consider *cocycles of isometries* of H, that is, cocycles $A \colon \Omega \to \mathrm{Isom}(H)$ over a dynamics $F \colon \Omega \to \Omega$, as in (1). We use the notation (2) for the products. The *maximal drift* of a cocycle of isometries is defined as

$$\mathrm{drift}(F, A) := \lim_{n\to\infty} \frac{1}{n} \sup_{\omega\in\Omega} d\big(A^{(n)}(\omega)p_0, p_0\big). \tag{10}$$

Notice that the limit exists by subadditivity, and is independent of the choice of $p_0 \in H$.

Remark 1.3. It follows from Kingman's subadditive ergodic theorem that for every ergodic probability measure μ for $F\colon \Omega \to \Omega$, the limit

$$\lim_{n\to\infty} \frac{1}{n} d\big(A^{(n)}(\omega)p_0, p_0\big)$$

exists for μ-almost every $\omega \in \Omega$, and is a constant (also independent of $p_0 \in H$). This is the *drift* of the cocycle of isometries with respect to the measure μ; let us denote it by $\mathrm{drift}(F, A, \mu)$. The following "variational principle" holds[4]:

$$\mathrm{drift}(F, A) = \sup_{\mu} \mathrm{drift}(F, A, \mu),$$

where μ runs over all invariant ergodic probabilities for F. $\triangleleft$

We say that the cocycle has *uniform sublinear drift* if $\mathrm{drift}(F, A) = 0$. By the Remark above, this happens if and only if A has zero drift with respect to every F-invariant probability measure.

The *displacement* of a (continuous) section $\varphi\colon \Omega \to H$ is defined by

$$\mathrm{displ}(\varphi) := \sup_{\omega\in\Omega} d\big(A(\omega)\varphi(\omega), \varphi(F\omega)\big) . \tag{11}$$

Notice that $\mathrm{displ}(\varphi) = 0$ if and only if the section φ is *invariant*, that is, $A(\omega)\varphi(\omega) = \varphi(F\omega)$ holds for every $\omega \in \Omega$.

It is not hard to show (see Proposition 4.3) that the displacement of any continuous section $\varphi\colon \Omega \to H$ is at least the maximal drift of the cocycle:

$$\mathrm{displ}(\varphi) \ge \mathrm{drift}(F, A). \tag{12}$$

Our first main result is a converse of (12):

Theorem A (Existence of sections of nearly minimal displacement; discrete time)**.** *Assume that H is a Busemann space. Given a cocycle $A\colon \Omega \to \mathrm{Isom}(H)$ over $F : \Omega \to \Omega$, for each $\varepsilon > 0$ there exists a continuous section $\varphi : \Omega \to H$ such that* $\mathrm{displ}(\varphi) \le \mathrm{drift}(F, A) + \varepsilon$.

Thus, together with (12), this theorem implies that the maximal drift is the infimum of the displacements of the continuous sections φ.

For the next result, we need extra assumptions on the space H, the most important being that H is a symmetric space (see § 2.5).

Let us say that a subset of $\Omega \times H$ is *bounded* if its projection on H is a bounded set.

Theorem B (Creating invariant sections; discrete time)**.** *Assume that H is*

- *either a proper*[5] *Busemann space;*

---

[4] This follows from [Schr, Thrm. 1] or [SS, Thrm. 1.7]. Although these references assume Ω to be compact metrizable, the proofs also work for compact Hausdorff Ω. (See also the proof of Proposition 1 in [AB].) A particular case was considered in [Fu].

[5] A metric space is called *proper* if bounded closed sets are compact.

- *or a space of bounded nonpositive curvature in the sense of Alexandrov*[6]*.*

Also assume that H is symmetric. If A is a cocycle of isometries of H with uniform sublinear drift, then there is a sequence of cocycles $A_N : \Omega \to H$ such that:

- *each A_N has a (continuous) invariant section;*
- *the sequence of skew-product maps $G_N(\omega, p) = (F(\omega), A_N(\omega)p)$ converges uniformly to $G(\omega, p) = (F(\omega), A(\omega)p)$ on bounded subsets of $\Omega \times H$.*

Remark 1.4. We can also show that the cocycles A_N given by Theorem B are all cohomologous to cocycles taking values in the stabilizer in $\mathrm{Isom}(H)$ of a point $p_0 \in H$. (In the case of the ellipsoids, this is isomorphic to $\mathrm{O}(d)$.) $\triangleleft$

Actually, the results above also hold in the a more general setting, namely when the trivial bundle $\Omega \times H$ is replaced by a fiber bundle whose fibers are isometric to H; see Section 4.

1.4 The general main results: continuous time

Now we assume that H is a Cartan–Hadamard manifold, that is, a complete simply connected Riemannian manifold (possibly of infinite dimension, modeled on a Hilbert space) of nonpositive sectional curvature. (See § 3.2 for details.)

A *semiflow* $\{F^t\}$ on Ω is a continuous-time dynamical system, that is, a continuous map $(\omega, t) \in \Omega \times \mathbb{R}_+ \mapsto F^t\omega \in \Omega$ such that $F^0 = \mathrm{id}$ and $F^{t+s} = F^t \circ F^s$ for all s, t in $\mathbb{R}$.

A *cocycle of isometries* (of H) over $\{F^t\}$ is a 1-parameter family of maps $A^{(t)} \colon \Omega \to \mathrm{Isom}(H)$ (where $t \in \mathbb{R}_+$) satisfying

$$A^{(0)}(\omega) = \mathrm{id}, \quad A^{(s+t)}(\omega) = A^{(s)}(F^t\omega)A^{(t)}(\omega) \tag{13}$$

and such that

$$A^{(t)}(\omega)p \text{ is continuous w.r.t. } (t, \omega, p) \text{ and continuously differentiable w.r.t. } t. \tag{14}$$

Associated to $\{A^t\}$ there is a vector-field $a(\omega)$ on H defined by

$$a(\omega)(p) = \frac{\partial}{\partial t} A^{(t)}(\omega)p \bigg|_{t=0} . \tag{15}$$

This defines a continuous map $a \colon \Omega \to \mathrm{Kill}(H)$, where $\mathrm{Kill}(H)$ is the set of Killing fields on H. (See § 3.2 for the topology on $\mathrm{Kill}(H)$.) Conversely, given any continuous map $a \colon \Omega \to \mathrm{Kill}(H)$, there is a unique cocycle of isometries $(F^t, A^{(t)})$ satisfying the ODE

$$\frac{\partial}{\partial t} A^{(t)}(\omega)p = a(F^t\omega)(A^{(t)}(\omega)p) . \tag{16}$$

---

[6] See § 2.6 for the definitions.

The map a is called the *infinitesimal generator* of the cocycle.

We define the *maximal drift* of a continuous-time cocycle of isometries in the same way as the discrete-time situation:

$$\operatorname{drift}(F, A) := \lim_{t \to \infty} \frac{1}{t} \sup_{\omega \in \Omega} d\big(A^{(t)}(\omega) p_0, p_0\big), \quad p_0 \in H \text{ arbitrary.} \tag{17}$$

Again, we say that the cocycle has *uniform sublinear drift* if $\operatorname{drift}(F, A) = 0$.

A (continuous) section $\varphi \colon \Omega \to H$ is said to be *differentiable with respect to the semiflow*[7] if for every $\omega \in \Omega$, the derivative

$$\varphi'(\omega) := \frac{\partial}{\partial t} \varphi(F^t \omega) \bigg|_{t=0} \tag{18}$$

exists and defines a continuous map $\varphi' \colon \Omega \to TH$. If φ is a section that is differentiable with respect $\{F^t\}$, then its *speed* with respect to $\{A^t\}$ is defined by:

$$\operatorname{speed}(\varphi) = \sup_{\omega \in \Omega} \left\| a(\omega)(\varphi(\omega)) - \varphi'(\omega) \right\|, \tag{19}$$

where a is the infinitesimal generator of the cocycle. The speed is the continuous-time analogue of the displacement (11). Notice that $\operatorname{speed}(\varphi) = 0$ holds if and only if the section φ is *invariant*, that is, $A^{(t)}(\omega)\varphi(\omega) = \varphi(F^t \omega)$.

Analogously to (12), we have

$$\operatorname{speed}(\varphi) \geq \operatorname{drift}(F, A). \tag{20}$$

The continuous-time versions of Theorems A and B are given below:

Theorem C (Existence of sections of nearly minimal speed; continuous time)**.** *Assume that H is a Cartan–Hadamard manifold. Given a continuous-time cocycle of isometries $\{A^{(t)}\}$ of H over a semiflow $\{F^t\}$ on Ω, for each $\varepsilon > 0$ there exists a continuous section $\varphi \colon \Omega \to H$ that is differentiable with respect to the semiflow $\{F^t\}$ and such that* $\operatorname{speed}(\varphi) \leq \operatorname{drift}(F, A) + \varepsilon$.

Theorem D (Creating invariant sections; continuous time)**.** *In the context of the preceding theorem, assume moreover that H is a symmetric space and that the cocycle $\{A^{(t)}\}$ has uniform sublinear drift. Then there is $\tilde{a}$ arbitrarily close to a such that the associated cocycle $\{\tilde{A}^{(t)}\}$ has an invariant continuous section.*

Here, the convergence of a sequence $\tilde{a}_N \colon \Omega \to \operatorname{Kill}(H)$ to a is in the following sense: for each bounded subset B of H, the sequence $\tilde{a}_N(\omega)(p)$ converges to $a(\omega)(p)$ uniformly with respect to $(\omega, p) \in B \times \Omega$. (See § 3.2 for more precise explanation about the topologies.)

---

[7] A similar definition appears in [Schw].

1.5 Invariant conformal structures for matrix cocycles

We return to matrix cocycles and describe another application of out results.

A *Riemannian metric* on the vector bundle $\Omega \times \mathbb{R}^d$ is a continuous choice of inner product $\langle \cdot, \cdot \rangle_\omega$ on each fiber $\{\omega\} \times \mathbb{R}^d$. We denote $\|v\|_\omega = \sqrt{\langle v, v \rangle_\omega}$.

Theorem 1.5. *Assume that* $F \colon \Omega \to \Omega$ *is a uniquely ergodic homeomorphism with an invariant measure of full support. Then for every cocycle* A *in a dense subset* $C^0(\Omega, \mathrm{GL}(d, \mathbb{R}))$ *there exist:*

- *a Riemannian metric on the vector bundle* $\Omega \times \mathbb{R}^d$*;*
- *a continuous* (F, A)*-invariant splitting* $\mathbb{R}^d = E_1(\omega) \oplus \cdots \oplus E_k(\omega)$ *which is orthogonal with respect to the Riemannian metric;*
- *constants* $c_1 > \cdots > c_k > 0$ *such that*

$$\|A(\omega) v_i\|_{T\omega} = c_i \|v_i\|_\omega \quad \text{for all } v_i \in E_i(\omega). \tag{21}$$

Theorem 1.5 follows by combining the fibered versions of the previous results together with a result from [BV]. The complete proof is given in Section 5, where we also present a version of the theorem for minimal homeomorphisms.

Remark 1.6. For another result that yields invariant conformal structures for linear cocycles (under different hypotheses), see [KalS]. ◁

1.6 Further remarks and questions

In the case of isometries of the line $\mathbb{R}$, there are "one-sided" versions of our results, as we now explain.

Let $g \colon \Omega \to \mathbb{R}$ be a continuous function, and consider the skew-product $G = G_{F,g}$ given by (8). We define the *maximal upward drift* as:

$$\mathrm{drift}^+(F, g) := \sup_\mu \int_\Omega g \, \mathrm{d}\mu \,, \tag{22}$$

where μ runs over invariant ergodic probabilities for F.[8]

If $\varphi \colon \Omega \to \mathbb{R}$ is a continuous section, we define its *upward displacement* (with respect to (F, g)) as:

$$\mathrm{displ}^+(\varphi) := \sup_\Omega (g + \varphi - \varphi \circ F) \,. \tag{23}$$

This is a measure of how much the graph of φ is moved upwards by the action of the skew-product G. Notice that $\mathrm{displ}^+(\varphi) \geq \mathrm{drift}^+(F, g)$ for every φ.

The one-sided analogue of Theorem A with $H = \mathbb{R}$ is the following:

Theorem 1.7. *For every* $\varepsilon > 0$ *there is a continuous function* $\varphi \colon \Omega \to \mathbb{R}$ *such that* $\mathrm{displ}^+(\varphi) \leq \mathrm{drift}^+(F, g) + \varepsilon$*.*

---

[8] The study of this quantity (and of the measures which realize the supremum) is called *ergodic optimization*; see [Je].

In other words, given any g, there is $\tilde{g}$ cohomologous to g such that $\tilde{g} \le \operatorname{drift}^+(F,g) + \varepsilon$. [9] For an application of this result for matrix cocycles, see Remark 5.2.

Proof of Theorem 1.7. [10] Let $a = \operatorname{drift}^+(F,g) + \varepsilon$ and let

$$\varphi(\omega) = \sup_{n \ge 0} \left(g^{(n)}(\omega) - an \right). \tag{24}$$

To see that this formula gives a well-defined continuous function, first notice (using a Krylov–Bogoliubov argument) that there exists $n_0 \in \mathbb{Z}_+$ such that $g^{(n)}(\omega)/n < a$ for every $n > n_0$ and every $\omega \in \Omega$. In particular, one only needs to consider $n \le n_0$ in the supremum (24); this shows that φ is continuous.

Moreover,

$$\varphi(F(\omega)) = \sup_{n \ge 1} \left(g^{(n)}(\omega) - an \right) - g(\omega) + a \le \varphi(\omega) - g(\omega) + a,$$

as desired. $\square$

We now discuss some questions that are suggested by our results.

The study of the probability measures μ which realize the supremum in (22) is an interesting problem with rich ramifications; see [Je]. So it could be also interesting to study *drift maximizing measures*, i.e., those which realize the supremum in (10).

All our results concern approximations in the C^0 topology. It is natural to ask whether the differentiability class can be improved. However, this is non-trivial already in the classical case (i.e., that of Theorem 1.2), where it is closely related to the existence of invariant distributions for the base dynamics. (See [Kat, AK].) We do not know if such a relation can be extended to the cocycles considered in this work.

It is also natural to look for conditions under which we can improve the conclusion of Theorem 1.1 and find a perturbed cocycle cohomologous to a constant rotation, or even to the identity. The case of $\mathrm{SL}(2,\mathbb{R})$-cocycles is studied in [ABD2]. We expect that progress in this direction would allow us to deal with the problem of genericity of projective hyperbolicity (i.e., existence of dominated splittings) in the space of linear cocycles.

A further problem is to consider linear cocycles (or even cocycles of isometries) over actions of more complicated semigroups than $\mathbb{Z}_+$ or $\mathbb{R}_+$. Again, this is a difficult problem even in the classical case, where the analogue of Theorem 1.2 is know to hold for nilpotent group actions (see [MOP2]) but not for solvable group actions (see [MOP1]). In spite of that, it seems hard trying to show analogous statements for cocycles of isometries over a nilpotent group, as there already are deep open problems about nilpotent group actions by isometries of CAT(0) spaces (see, for instance, [ANT, Pau]).

---

[9] In general it is not possible to solve to solve this problem for $\varepsilon = 0$ (for example, take F a Liouville rotation of the circle and g a suitable function). Nevertheless, this can be done under some hyperbolicity and regularity assumptions; see [CLT, Prop. 11], [Je, Thrm. 4.7].

[10] A similar trick appears in the proof of [CLT, Prop. 11]. The result can also be proved less directly by using the Fenchel–Rockafellar duality theorem; see [GL, Thrm. 1].

1.7 Organization of the paper

Although the main ideas of the proofs of the discrete-time results and the continuous-time results are similar, the technical details are somewhat different. Thus we prove two kinds of results in a nearly independent way.

In Section 2 we prove Theorems A, B and 1.1, while in Section 3 we prove Theorems C and D. Along the way, we explain the geometrical tools and properties that are required for these proofs. Some of these properties become simpler to prove if the space H is proper (i.e., finite-dimensional in the manifold case). Thus in Sections 2 and 3, we give proofs for the proper case, and leave for Appendices A and B the proofs of those lemmas in the general case. A technical property which is needed for the proof of Theorem D is proved in Appendix C. In Section 4, we prove fibered versions of Theorems A and B, and in Section 5, we apply these to prove Theorem 1.5.

Acknowledgements. We thank A. Avila, E. Garibaldi, N. Gourmelon, A. Karlsson, A. Kocsard, M. Ponce and R. Tessera for valuable discussions. We also thank the hospitality of the Institut de Mathématiques de Bourgogne, where this work started.

2 Proof of the results for discrete-time cocycles

2.1 Busemann spaces

Let (H, d) be a separable metric space. We say that H is a *geodesic space* if it is complete and every two points p, q in H can joined by a geodesic, that is, a curve $\gamma\colon [0,1] \to H$ such that $\gamma(0) = p$, $\gamma(1) = q$, and $d(\gamma(t), \gamma(s)) = |t - s| d(p, q)$ for all s, t in $[0, 1]$. If these curves are unique for arbitrarily prescribed p, q, then we say that H is *uniquely geodesic*.

The space H has *nonpositive curvature in the sense of Busemann* (it is a *Busemann space*, for short) if it geodesic and the distance function along geodesics is convex. Equivalently, given any two pairs of points p, q and p', q', their corresponding midpoints m, m' satisfy

$$d(m, m') \leq \frac{d(p, p')}{2} + \frac{d(q, q')}{2}. \tag{25}$$

The family of such spaces obviously includes all strictly convex Banach spaces. (General Banach spaces may also be included in this category when considering only segments of lines as geodesics.) It also includes simply connected Riemannian manifolds of nonpositive curvature, as for example the space of ellipsoids.

(See §2.9 for more details as well as the discussion of infinite-dimensional extensions of this example.)

2.2 Barycenter maps

Given a metric space (H, d), we denote by $\mathcal{P}^1(H)$ the space of probability measures on H with finite first moment, that is, such that for some (equivalently, for every) $p_0 \in H$ one has

$$\int_H d(p_0, p)\,\mathrm{d}\mu(p) < \infty.$$

We endow this space with the 1-Wasserstein metric W_1 defined as

$$W_1(\mu, \nu) := \inf_{P \in (\mu|\nu)} \int_{H \times H} d(x, y)\,\mathrm{d}P(p, q), \tag{26}$$

where $(\mu \mid \nu)$ denotes the set of all *couplings* of μ and ν, that is, all probability measures P on $H \times H$ whose projection along the first (resp. second) coordinate coincides with μ (resp. ν).

Example 2.1. If μ, ν_1, $\nu_2 \in \mathcal{P}^1(H)$ then for every $\lambda \in [0, 1]$,

$$W_1\big((1-\lambda)\mu + \lambda\nu_1, (1-\lambda)\mu + \lambda\nu_2\big) \le \lambda \cdot \sup\{d(p_1, p_2);\ p_i \in \operatorname{supp} \nu_i\}.$$

Indeed, this follows directly from the definitions by considering the coupling $(1-\lambda)i_*\mu + \lambda\,\nu_1 \times \nu_2$, where $i(p) := (p, p)$.

For much more on Wasserstein metrics, see e.g. [Vi].

Theorem 2.2. *If H is a Busemann space, then there exists a map* $\operatorname{bar}\colon \mathcal{P}^1(H) \to H$ *that is equivariant with respect to the action of the isometries and* 1*-Lipschitz for the* 1*-Wasserstein metric on* $\mathcal{P}^1(H)$.

In particular, in the situation of Example 2.1

$$d\big(\operatorname{bar}((1-\lambda)\mu + \lambda\nu_1), \operatorname{bar}((1-\lambda)\mu + \lambda\nu_2)\big) \le \lambda \cdot \sup\{d(p_1, p_2);\ p_i \in \operatorname{supp} \nu_i\}. \tag{27}$$

We also denote

$$\operatorname{bar}(p_1, \ldots, p_n) := \operatorname{bar} \frac{1}{n}(\delta_{p_1} + \cdots + \delta_{p_n}),$$

where δ_p denotes Dirac measure on p. So it follows from (27) that

$$d\big(\operatorname{bar}(p_1, \ldots, p_{n-1}, p_n), \operatorname{bar}(p_1, \ldots, p_{n-1}, p'_n)\big) \le \frac{1}{n} d(p_n, p'_n). \tag{28}$$

Henceforth, we will not need to know what precisely is the map bar above, although its geometrical flavor should be intuitively transparent. For instance, $\operatorname{bar}(p)$ coincides with p, while $\operatorname{bar}(p_1, p_2)$ is the midpoint between p_1 and p_2. The definition of $\operatorname{bar}(p_1, p_2, p_3)$ is, however, quite involved.

In its full generality, the theorem above was proved by Navas in [Na] by elaborating on an idea introduced by Es-Sahib and Heinich in [EH]. Nevertheless, for compactly supported measures μ on CAT(0)-spaces, a much more classical notion of barycenter due to Cartan (see [Ca, p. 267]) is enough for our purposes. (See § 2.6 for the definition of CAT(0) spaces; see also [BK, Jo, AL].) The Cartan barycenter of μ as above is defined as the unique point that minimizes the function

$$f_\mu(p) = \int_H d^2(q,p)\, \mathrm{d}\mu(q). \tag{29}$$

The fact that Cartan's barycenter is 1-Lipschitz with respect to the 1-Wasserstein metric is proved in [St].

2.3 Existence of sections of nearly minimal displacement: proof of Theorem A

Proof of Theorem A. Given a cocycle A of isometries of a Busemann space H over $F\colon \Omega \to \Omega$, we fix any $p_0 \in H$. We introduce a formula analogous to (7):

$$\varphi_N(\omega) := \operatorname{bar}\left(p_0, A(\omega)^{-1}p_0, [A^{(2)}(\omega)]^{-1}p_0, \dots, [A^{(N-1)}(\omega)]^{-1}p_0\right), \tag{30}$$

where bar stands for the barycenter introduced in Theorem 2.2. Then, by equivariance of the barycenter, we have (see Fig. 1):

$$\begin{aligned} A(\omega)\varphi_N(\omega) &= \operatorname{bar}\left(A(\omega)p_0, p_0, A(F\omega)^{-1}p_0, \dots, [A^{(N-2)}(F\omega)]^{-1}p_0 \right), \\ \varphi_N(F\omega) &= \operatorname{bar}\left(p_0, A(F\omega)^{-1}p_0, \dots, [A^{(N-2)}(F\omega)]^{-1}p_0, [A^{(N-1)}(F\omega)]^{-1}p_0\right). \end{aligned}$$

Using (28), we have

$$\begin{aligned} d\big(A(\omega)\varphi_N(\omega), \varphi_N(F\omega)\big) &\le \frac{1}{N} d\big(A(\omega)p_0, [A^{(N-1)}(F\omega)]^{-1}p_0\big) \\ &= \frac{1}{N} d\big(A^{(N)}(\omega)p_0, p_0\big). \end{aligned}$$

In particular,

$$\operatorname{displ}(\varphi_N) \le \frac{1}{N} \sup_{\omega\in\Omega} d\big(A^{(N)}(\omega)p_0, p_0\big).$$

The theorem then follows by taking $\varphi = \varphi_N$ with sufficiently large N (depending on ε). ☐

Remark 2.3. The same method of proof yieds the following: There exists a family $\{\varphi_\tau\}_{\tau\in\mathbb{R}_+}$ of continuous sections such that $\lim_{\tau\to\infty} \operatorname{displ}(\varphi_\tau) = \operatorname{drift}(F,A)$; moreover φ_τ depends continuously on τ and also on A. (Such parameterized sections play an important role in [ABD2].) Furthermore, such association is continuous: given a continuous family $A^{(s)}$ of cocycles (s lives in an arbitrary topological space), the resulting $\varphi_t^{(s)}$ are jointly continuous. $\triangleleft$



Figure 1: Checking almost invariance of the section φ_N.

2.4 Isometries

Recall that the *displacement function* of $J \in \mathrm{Isom}(H)$ is the function

$$p \in H \mapsto d(J(p), p). \tag{31}$$

Since H is a Busemann space, this function is convex. The infimum of the displacement function is called the *translation length* of J.

On the space $\mathrm{Isom}(H)$, we consider the first-countable topology for which the convergence of sequences is uniform convergence on bounded subsets. This is called the *bounded-open topology.*[11]

We also endow the set $C(\Omega, \mathrm{Isom}(H))$ of continuous functions from Ω into $\mathrm{Isom}(H)$ with the compact-open topology. A sequence (A_n) converges to A in this topology if and only if for every bounded subset B of H, the sequence $(A_n(\omega)(p))$ converges to $A(\omega)(p)$ uniformly with respect to $(\omega, p) \in \Omega \times B$.

2.5 Symmetric geodesic spaces, transvections, and a displacement estimate

We say that a uniquely geodesic space H is *geodesically complete* if the maximal interval of definition of all geodesics is $\mathbb{R}$. For such a space, the *symmetry* at a point $p_0 \in H$ is the map $\sigma_{p_0} \colon H \to H$ that sends p to to the unique point p' such that p_0 is the midpoint between p and p'. So σ_{p_0} is an involution. We say that H is a *symmetric geodesic space* if σ_{p_0} is a isometry for every $p_0 \in H$, and the map $(p_0, p) \mapsto \sigma_{p_0}(p)$ is continuous.

---

[11] If H is proper then this coincides with the compact-open topology, which is the usual topology on $\mathrm{Isom}(H)$; see e.g. [He].

Assume that H is a symmetric geodesic space. Following É. Cartan, we call a *transvection* an isometry of the form $J = \sigma_{p_2} \circ \sigma_{p_1}$. If $\gamma : \mathbb{R} \to H$ is a isometric (unit-speed) parametrization of the geodesic passing through p_1 and p_2, say with $\gamma^{-1}(p_1) < \gamma^{-1}(p_2)$, then $J(\gamma(t)) = \gamma(t+b)$ holds for all $t \in \mathbb{R}$, where $b = 2d(p_1, p_2)$. We say that J translates the geodesic γ by length b.

We remark that if, in addition, H is a Busemann space, then $d(J(q), q) \geq b$ for all $q \in H$; see [Pa]. So the translation length of J is precisely b.

Lemma 2.4. *Assume that H is a symmetric geodesic space. Also assume that H is proper. Let J be a transvection that translates a geodesic γ by length b. Then*

$$d(J(q), q) \leq f\left(b, d(q, \gamma)\right) \text{ for every } q \in H, \tag{32}$$

where $f \colon \mathbb{R}_+ \times \mathbb{R}_+ \to \mathbb{R}_+$ is a function that depends only on the space H, and is monotonically increasing with respect to each variable.

Proof. Define a function $\tilde{f} \colon H \times \mathbb{R}_+ \times \mathbb{R}_+ \to \mathbb{R}_+$ by

$$\tilde{f}(p_0, b, \ell) = \sup\left\{d(J(q), q);\ J = \sigma_{p_1} \circ \sigma_{p_0},\ d(p_1, p_0) \leq b/2,\ d(q, p_0) \leq \ell\right\}.$$

The supremum is finite by properness of H and continuity. Since the group of isometries acts transitively on H, the value $\tilde{f}(p_0, b, \ell)$ actually does not depend on p_0; call it $f(b, \ell)$. Then (32) holds. ☐

Remark 2.5. It follows that under the assumptions of Lemma 2.4, the map $p_0 \in H \mapsto \sigma_{p_0} \in \mathrm{Isom}(H)$ is continuous (where $\mathrm{Isom}(H)$ is endowed with the bounded-open topology, as explained in § 2.4.) ◁

2.6 Curvature bounds in the sense of Alexandrov, and a displacement estimate

In the case that H is infinite dimensional, the proof of Lemma 2.4 given above obviously doesn't work. Nevertheless, the lemma holds if properness is replaced by some curvature hypotheses, as we next explain. Readers who are not interested in infinite dimensional applications can skip this subsection.

Let H be a uniquely geodesic space. A *triangle* $\triangle(p_1, p_2, p_3)$ in H consists on three points p_1, p_2, p_3 and three geodesic segments joining them. Given $\kappa \leq 0$, the *model space* (M_κ, d_κ) is the two-dimensional plane of constant curvature κ. If H is uniquely geodesic, then we say that H has *curvature $\leq \kappa$ (resp. $\geq \kappa$) in the sense of Alexandrov* if for every triangle $\triangle(p_1, p_2, p_3)$ in H, the following inequality holds for all $t \in [0, 1]$:

$$d\big(p_3, (1-t)p_1 + tp_2\big) \leq \ \text{(resp. } \geq \text{)}\ d_\kappa\big(\tilde{p}_3, (1-t)\tilde{p}_1 + t\tilde{p}_2\big),$$

where $\triangle(\tilde{p}_1, \tilde{p}_2, \tilde{p}_3)$ is a triangle in M_κ such that $d(p_i, p_j) = d_\kappa(\tilde{p}_i, \tilde{p}_j)$ for all i, j in $\{1, 2, 3\}$. We say that $\triangle(\tilde{p}_1, \tilde{p}_2, \tilde{p}_3)$ is a *SSS-comparison triangle*[12] for

---

[12] SSS stands for *side-side-side*.

$\triangle(p_1, p_2, p_3)$. Spaces of curvature ≤ 0 are also called *CAT(0)-spaces*. It is a standard fact that every complete CAT(0) space is a Busemann space (see [BH, p. 176] or [St, Corollary 2.5]).

We will say that an uniquely geodesic space H has *bounded nonpositive curvature in the sense of Alexandrov* if it has curvature ≤ 0 and $\geq \kappa$ for some $\kappa \leq 0$.

Now we have the following version of Lemma 2.4:

Lemma 2.6. *Assume that H is a symmetric space of bounded nonpositive curvature in the sense of Alexandrov. Let J be a transvection that translates a geodesic γ by length b. Then*

$$d(J(q), q) \leq f\left(b, d(q, \gamma)\right) \text{ for every } q \in H, \tag{33}$$

where $f\colon \mathbb{R}_+ \times \mathbb{R}_+ \to \mathbb{R}_+$ is a function that depends only on the space H, and is monotonically increasing with respect to each variable.

We leave the proof of this lemma to Appendix A.

Remark 2.7. Similarly to Remark 2.5, we conclude that under the assumptions of Lemma 2.6, the map $p_0 \in H \mapsto \sigma_{p_0} \in \mathrm{Isom}(H)$ is continuous. $\triangleleft$

2.7 Macroscopic uniform homogeneity

Every symmetric space is homogeneous (in the sense that the group of isometries acts transitively). We will need however a stronger property, given by the following lemma:

Lemma 2.8 (Macroscopic uniform homogeneity)**.** *Assume that H is*

- *either a proper Busemann space;*
- *or a space of bounded nonpositive curvature in the sense of Alexandrov.*

Also assume that H is symmetric. Then there exists a continuous map $J\colon H \times H \to \mathrm{Isom}(H)$ with the following properties:

1. *$J(p, q)p = q$ for all p, q in H.*
2. *$J(p, q)$ converges to the identity as the distance between p and q converges to zero.*

More explicitly, assertion 2 means that for every $\varepsilon > 0$ and each bounded subset $B \subset H$ there exists $\delta > 0$ such that $d(J(p, q)r, r) < \varepsilon$ holds for all $r \in B$ whenever $d(p, q) < \delta$. (Notice that p and q are not restricted to a bounded set.)

Proof. Fix some $p_0 \in H$, and consider the transvection (see Fig. 2):

$$J(p, q) := \sigma_m \circ \sigma_{p_0}\,, \quad \text{where } m \text{ is the midpoint between } \sigma_{p_0}(p) \text{ and } q. \tag{34}$$

Applying the Busemann inequality (25) to the points $\sigma_{p_0}(p)$, p and q, we obtain $d(p_0, m) \leq \frac{1}{2} d(p, q)$. Therefore, the length by which the transvection $J(p, q)$ translates γ is at most $d(p, q)$. So assertion 2 follows directly from Lemma 2.4 or Lemma 2.6, according to the case. $\square$



Figure 2: m is the midpoint between q and $\sigma_{p_0}(p)$. The isometry $J(p,q) := \sigma_m \circ \sigma_{p_0}$ sends p to q and translates the geodesic joining p_0 and m by length $2d(p_0,m) \le d(p,q)$.

Remark 2.9. In the case where H is the hyperbolic plane, Lemma 2.8 follows from Lemma 5 from [ABD1]. Actually, our formula (34) reproduces the construction of that paper in this particular case. ◁

2.8 Creating invariant sections: proof of Theorem B

Proof of Theorem B. By Theorem A, there exists a sequence of sections φ_N such that $\lim_{N\to\infty} \operatorname{displ}(\varphi_N) = 0$. Define

$$\tilde{A}_N(\omega) = J\big(A(\omega)\varphi_N(\omega), \varphi_N(F\omega)\big) \circ A(\omega), \tag{35}$$

where J is given by Lemma 2.8. Then:

- $\tilde{A}_N(\omega)\varphi_N(\omega) = \varphi_N(F\omega)$, that is, φ_N is $\tilde{A}_N$-invariant.
- for each bounded subset B of H, the sequence $\tilde{A}_N(\omega)p$ converges to $A(\omega)p$ uniformly with respect to $(\omega,p) \in B \times \Omega$.

Thus the theorem is proved. □

Let us notice some extra properties of the invariant sections we have just constructed.

Remark 2.10. Take any point $p_0 \in H$ and consider $B_N(\omega) = U(F\omega)^{-1} \circ A_N(\omega) \circ U(\omega)$, where $U(\omega) = J(p_0, \varphi_N(\omega))$. Then B_N is a cocycle cohomologous to A_N that takes values in the stabilizer of p_0. This justifies Remark 1.4. ◁

Remark 2.11. Let us see how to obtain certain "accessibility" properties, which play an important role in [ABD2].

First, we extend the family $\{\varphi_N\}_{N\in\mathbb{Z}_+}$ to a family $\{\varphi_t\}_{t\in\mathbb{R}}$ by letting

$$\varphi_t(\omega) := \operatorname{bar}\left((1-t+N)\delta_{\varphi_N(\omega)} + (t-N)\delta_{\varphi_{N+1}(\omega)}\right), \quad \text{where } N = \lfloor t \rfloor.$$

Then $\operatorname{displ}\varphi_t \to 0$ as $t \to +\infty$. Repeating the construction of the proof of Theorem B, we conclude the following: *For every cocycle A with uniform sublinear drift, there exists a continuous family of cocycles A_t, $t \in [0,\infty]$, satisfying $A_\infty = A$ and such that for each $t < \infty$, A_t has a continuous invariant section φ_t (that also depends continuously on t). Moreover, such correspondence is continuous: given a continuous family $A(s)$ of cocycles (s in an arbitrary topological space), the resulting $A_t(s)$ and $\varphi_t(s)$ are jointly continuous.* ◁

2.9 Proof of Theorem 1.1

Fix an integer $d \geq 2$. The space $\mathcal{E}_d$ of ellipsoids in $\mathbb{R}^d$ centered at the origin can be identified with the space Pos_d of symmetric positive definite $d\times d$ matrices; to each such matrix P, we can associate the ellipsoid $\{u \in \mathbb{R}^d;\ \langle P^{-1}u, u\rangle_{\mathrm{eucl}} \leq 1\}$. Notice that Pos_d also identifies with the cosets space $\mathrm{GL}(d,\mathbb{R})/\mathrm{O}(d)$ by writing each P above in the form $P = xx^T$, with $x \in \mathrm{GL}(d,\mathbb{R})$.

Let us recall some facts about the space Pos_d; see [Lan, Chapter XII] for details. The vector space Sym_d of symmetric $d \times d$ matrices identifies with the Lie algebra of Pos_d. Moreover, we have $\mathrm{Pos}_d = \exp(\mathrm{Sym}_d)$, where exp is the usual exponential map for matrices. Consider the following inner product on Sym_d:

$$\langle v, w\rangle_{\mathrm{id}} = \mathrm{tr}(vw).$$

(The reason of the id subscript will be clear soon.) Using this inner product, we may define a Riemannian structure on Pos_d. We identify the tangent space of Pos_d at a point $x \in \mathrm{Pos}_d$ with Sym_d (just regarding Pos_d as an open subset of the vector space Sym_d), and for each v, w in $T_x\mathrm{Pos}_d \sim \mathrm{Sym}_d$ we set

$$\langle v, w\rangle_x = \langle x^{-1}v, x^{-1}w\rangle_{\mathrm{id}}.$$

Endowed with this metric, the space Pos_d has nonpositive curvature. Moreover, the group $\mathbb{G} = \mathrm{GL}(d,\mathbb{R})$ acts by isometries on Pos_d as follows:

$$g : x \mapsto g(x) := g \cdot x \cdot g^T\ ,$$

where g^T is the transpose of the matrix g. (In terms of the identification of Pos_d with $\mathcal{E}_d$, this corresponds to taking the image of the ellipsoids under g.) In fact, every isometry of Pos_d is of this form. The stabilizer of the identity matrix is the orthogonal group $\mathrm{O}(d)$.

Finally, let us recall a formula for the distance between two points x, y in Pos_d. Namely, letting $\lambda_1, \dots, \lambda_d$ be the roots of $\det(ty - x) = 0$, we have

$$d_{\mathrm{Pos}_d}(x, y) = \sum_{i=1}^{d} (\log \lambda_i)^2. \tag{36}$$

In particular, the distance from $x \in \mathrm{Pos}_d$ to the identity equals the sum of the squares of the logarithms of its eigenvalues.

Proof of Theorem 1.1. Due to formula Eq. (36), a matrix cocycle has uniform subexponential growth if and only if the associated cocycle of isometries of Pos_d has uniform sublinear drift. Therefore, Theorem 1.1 follows from Theorem B. $\square$

Remark 2.12. Using [BH, Theorem. 10.58], we see that Theorem 1.1 can be extended to other groups of matrices, as complex matrices, quaternionic matrices, and symplectic matrices (in each case, the cohomologous cocycle after perturbation takes values in the corresponding maximal compact subgroup). The main ingredient is that these groups of matrices act by isometries on certain CAT(0) spaces, namely, the associated symmetric spaces. $\triangleleft$

Remark 2.13. Let $\mathcal{H}$ be infinite-dimensional separable real Hilbert space. Let $\mathrm{GL}^2(\infty, \mathbb{R})$ be the group of all invertible operators on $\mathcal{H}$ that may be written in the form $\mathrm{Id} + L$, where L is a Hilbert–Schmidt operator. Let $\mathrm{O}^2(\infty)$ be the orthogonal subgroup of $\mathrm{GL}^2(\infty, \mathbb{R})$. Then $H := \mathrm{GL}^2(\infty, \mathbb{R})/\mathrm{O}^2(\infty)$ is a symmetric Cartan–Hadamard manifold on which $\mathrm{GL}^2(\infty, \mathbb{R})$ acts by isometries; see [Lar]. Hence, Theorems A and B apply to this space. In particular, Theorem 1.1 extends to cocycles of this kind of operators.

Consider now the space $\mathrm{GL}(\infty, \mathbb{R})/\mathrm{O}(\infty)$, where $\mathrm{GL}(\infty, \mathbb{R})$ is the group of all bounded invertible operators on $\mathcal{H}$, and $\mathrm{O}(\infty)$ is the orthogonal subgroup. This is a Busemann space (see [CPR]), so Theorem A can be applied to $\mathrm{GL}(\infty, \mathbb{R})$-cocycles. Although this is also a symmetric space (as well as a Banach-manifold), we don't know if Theorem B applies. $\triangleleft$

3 Proof of the results for continuous-time cocycles

3.1 Cartan–Hadamard manifolds

Assume H is a Hilbert-manifold, that is, a separable C^∞-manifold modeled on a separable real Hilbert space $(\mathcal{H}, \langle \cdot, \cdot \rangle)$. Fix a Riemannian metric on H. (See [Lan] for the precise definition.)

If H is complete, simply connected, and has nonpositive sectional curvature, then H is called a *Cartan–Hadamard manifold*. In this case, the Cartan–Hadamard–McAlpin Theorem (see [Lan, § IX.3]) states that for each point $p \in H$, the exponential map $\exp_p \colon T_pH \to H$ is a diffeomorphism.

3.2 Killing fields and symmetric Cartan–Hadamard manifolds

Here we recall some general facts about symmetric manifolds and Killing fields; more information can be found in [Lan, Ch. XIII].

If H is Hilbert-manifold, a *Killing field* is a vector field that generates a (globally defined) flow of isometries. Then the flow also preserves the Riemannian connection. On the space $\mathrm{Kill}(H)$ of these fields, we consider the first-countable topology for which the convergence of sequences is uniform convergence on bounded subsets.

We endow $C(\Omega, \mathrm{Kill}(H))$ with the compact-open topology. Then a sequence (a_n) in $C(\Omega, \mathrm{Kill}(H))$ converges to a iff for every bounded set $B \subset H$, $\|a_n(\omega)(p) - a(\omega)(p)\|$ converges to 0 uniformly with respect to $(\omega, p) \in \Omega \times B$.

Now let H be a Cartan–Hadamard manifold. Suppose it is symmetric in the sense of § 2.5.[13] If $v_0 \in T_{p_0}H$ is a nonzero vector, let $\alpha\colon \mathbb{R} \to H$ be the geodesic such that $\alpha(0) = p_0$, $\alpha'(0) = v_0$. Consider the transvection

$$\tau_{\alpha,s} = \sigma_{\alpha(s/2)} \circ \sigma_{\alpha(0)}.$$

Then $\tau_{\alpha,s}$ is a flow of isometries called the *translation along* α. More precisely, we have $\tau_{\alpha,s}(\alpha(t)) = \alpha(t+s)$. Moreover, the derivative

$$T_{\alpha(t)}\tau_{\alpha,s}\colon T_{\alpha(t)}H \to T_{\alpha(t+s)}H$$

is the parallel transport along the geodesic α. Let ξ_{v_0} denote the Killing field that generates the flow $\tau_{\alpha,s}$. (For $v_0 = 0$, we define $\xi_{v_0} \equiv 0$.) Then the map $v_0 \in TH \mapsto \xi_{v_0} \in \mathrm{Kill}(H)$ is continuous.

As it is customary, we denote by $\mathfrak{m}_{p_0}$ the set of Killing fields ξ_{v_0}, where $v_0 \in T_{p_0}H$. This is a vector space, and it can also be expressed as

$$\mathfrak{m}_{p_0} = \big\{\xi \in \mathrm{Kill}(H);\ \nabla_\zeta \xi(p) = 0 \text{ for all vector fields } \zeta\big\},$$

where ∇ denotes covariant derivative.

3.3 Infinitesimal displacement estimates

The following lemma is the infinitesimal counterpart of Lemmas 2.4 and 2.6:

Lemma 3.1. *Let H be a symmetric Cartan–Hadamard manifold. There is a non-decreasing function $f\colon \mathbb{R}_+ \to \mathbb{R}_+$ with $f(0) = 1$ such that*

$$\|v_0\| \overset{\text{(I)}}{\leq} \|\xi_{v_0}(p)\| \overset{\text{(II)}}{\leq} f(d(p,p_0))\|v_0\|$$

for all p_0, $p \in H$, $v_0 \in T_{p_0}H$.

Inequality (I) above is related to nonpositive curvature; let us prove it first:

Proof of part (I) in Lemma 3.1. Fix p_0, $p \in H$, $v_0 \in T_{p_0}H$. Assume $v_0 \neq 0$, otherwise there is nothing to prove. Let β be a unit-speed geodesic joining p_0 and p. Let $\eta(t) := \xi_{v_0}(\beta(t))$. Then (see [Lan, Prop. 2.2, Ch XIII]) η is a Jacobi field over the geodesic β. By [Lan, Prop. 5.6, Ch. XIII], we have $\nabla_{\beta'}\eta(0) = 0$.

---

[13] It is easy to check that this agrees with the definition from [Lan, p. 359].

Let $g(t) := \|\eta(t)\|^2$. By nonpositive curvature, this function is convex: see [Lan, Lemma 1.1, Ch. X]. The same lemma also says that $g' = 2\langle \nabla_{\beta'}\eta, \eta\rangle$, which vanishes at $t = 0$. It follows that $g(t) \geq g(0)$ for all $t \in \mathbb{R}$. In particular, $\|\xi_{v_0}(p)\|^2 \geq \|\xi_{v_0}(p_0)\|^2 = \|v_0\|^2$, thus completing the proof of inequality (I). $\square$

If H is finite-dimensional then the existence of a function with property (II) in Lemma 3.1 is nearly trivial, and does not rely on nonpositive curvature:

Proof of part (II) in Lemma 3.1 assuming $\dim H < \infty$*.* Consider

$$\tilde{f}(p_0, \ell) := \sup\big\{\|\xi_{v_0}(p)\|;\ p \in H \text{ with } d(p, p_0) \leq \ell,\ v_0 \in T_{p_0}H \text{ with } \|v_0\| = 1\big\},$$

which is finite by compactness. Since $\mathrm{Isom}(H)$ acts transitively on H, the value $\tilde{f}(p_0, \ell)$ actually does not depend on p_0, and so defines a function $f(p_0)$ with the required properties. $\square$

The proof of (II) in the infinite-dimensional case requires geometrical arguments and is given in the Appendix B.

3.4 Infinitesimal uniform homogeneity

The following is an infinitesimal version of the macroscopic uniform homogeneity, i.e., Lemma 2.8. It basically says that we can move any point p in any desired direction w by an infinitesimal isometry (Killing field), and these fields can be chosen so that they converge uniformly (with respect to p) in bounded sets to zero as $\|w\| \to 0$.

Lemma 3.2 (Infinitesimal uniform homogeneity)**.** *Let $p_0 \in H$. There is a continuous map*

$$\begin{aligned} K\colon TH &\to \mathrm{Kill}(H) \\ w &\mapsto K_w \end{aligned}$$

with the following properties:

1. *The vector field K_w extends w, that is, if $p = \pi(w) \in H$ is the base point of w, then $K_w(p) = w$;*
2. *For any $q \in H$,*
$$\|K_w(q)\| \leq f(d(q, p_0))\|w\|,$$
where f is given by Lemma 3.1.

Proof of Lemma 3.2. Let $p_0 \in H$ be fixed. For $p \in H$, consider the map

$$\begin{aligned} L_p\colon T_{p_0}H &\to T_pH \\ v_0 &\mapsto \xi_{v_0}(p). \end{aligned}$$

We list below some properties of L_p:

- It is linear: see [Lan, p. 363].
- It is continuous: see part (II) of Lemma 3.1.
- It is one-to-one, and the inverse (on the image) is continuous: see part (I) of Lemma 3.1.
- It is onto: see Lemma B.1 in Appendix B. (In finite dimension, this part would of course be a trivial consequence of the others.)

Given $w \in T_pH$, define $K_w := \xi_{(L_p)^{-1}(w)}$. Using parts (II) and (I) of Lemma 3.1, we have

$$\begin{aligned} \|K_w(q)\| &\le f(d(q,p_0))\|K_w(p_0)\| \\ &\le f(d(q,p_0))\|w\|. \end{aligned} \qquad \square$$

Remark 3.3. It is possible (but with no advantage) to give an alternative proof Lemma 2.8 using Lemma 3.2: Given two points p and q, join them by a geodesic $\gamma\colon [0,\ell] \to H$, and integrate the time-varying Killing field $K_{\gamma'(t)}$ (where K is given by Lemma 3.2) to get the map $J(p,q)$. (In fact, the first proof of Lemma 5 from [ABD1] followed a similar idea, but this proof wasn't published because a more direct proof was found.) $\triangleleft$

3.5 Existence of sections of nearly minimal speed: proof of Theorem C

For each $p \in H$, we let δ_p or $\delta(p)$ denote the Dirac measure at the point p. If $\gamma\colon [a,b] \to H$ is a curve, we denote by $\int_a^b \delta(\gamma(t))\,\mathrm{d}t$ the measure on H obtained by pushing-forward by γ the Lebesgue measure on $[a,b]$.

In the proof of Theorem C, we will need the following technical result, whose proof is given in the Appendix C.

Lemma 3.4 (Differentiability of the Cartan barycenter)**.** *Let $I \subset \mathbb{R}$ be an open interval and let $h\colon I \times [0,T] \to H$ be a continuous mapping that is continuously differentiable with respect to the first variable. Then the map $\bar{h}\colon I \to H$ defined by*

$$\bar{h}(t) = \operatorname{bar}\left(\frac{1}{T}\int_0^T \delta_{h(t,s)}\,\mathrm{d}s\right) \tag{37}$$

(where bar *denotes the Cartan barycenter) is continuously differentiable.*

Proof of Theorem C. Fix any $p_0 \in H$. For $T > 0$, let

$$\varphi_T(\omega) = \operatorname{bar}\left(\frac{1}{T}\int_0^T \delta\left([A^{(t)}(\omega)]^{-1}p_0\right)\,\mathrm{d}t\right) \tag{38}$$

(Compare with (30).)

Claim. *The function $\varphi_T\colon \Omega \to H$ is differentiable with respect to the semiflow.*

Proof of the claim. We have

$$\varphi_T(F^t\omega) = \operatorname{bar}\left(\frac{1}{T}\int_0^T \delta\left([A^{(s)}(F^t\omega)]^{-1}p_0\right)\,\mathrm{d}s\right).$$

So, by Lemma 3.4, in order to check that the map $\varphi_T(F^t\omega)$ is continuously differentiable with respect to t, it is sufficient to check that the map $(s,t)\mapsto [A^{(s)}(F^t\omega)]^{-1}p_0 \in H$ is continuous and continuously differentiable with respect to t. But these properties follow from the cocycle identity (13) and the regularity assumptions (14). $\square$

Next, we want to estimate the distance:

$$d\big(\varphi_T(F^t\omega), A^{(t)}(\omega)\varphi_T(\omega)\big) = d\big(\underbrace{[A^{(t)}(\omega)]^{-1}(\varphi_T(F^t\omega))}_{(\star)}, \varphi_T(\omega)\big). \tag{39}$$

Assuming $t\in(0,T)$, we have

$$\begin{aligned}(\star) &= \operatorname{bar}\frac{1}{T}\int_0^T \delta\left([A^{(t)}(\omega)]^{-1}[A^{(s)}(F^t\omega)]^{-1}p_0\right)\,\mathrm{d}s\\ &= \operatorname{bar}\frac{1}{T}\int_0^T \delta\left([A^{(s+t)}(\omega)]^{-1}p_0\right)\,\mathrm{d}s\\ &= \operatorname{bar}\frac{1}{T}\int_t^{T+t} \delta\left([A^{(s)}(\omega)]^{-1}p_0\right)\,\mathrm{d}s.\end{aligned}$$

Using the barycenter property (27), we obtain that the distance (39) is at most

$$\frac{t}{T}\sup\left\{d\big([A^{(s)}(\omega)]^{-1}p_0, [A^{(u)}(\omega)]^{-1}p_0\big);\ s\in[0,t], u\in[T,T+t]\right\}.$$

Dividing by t and making $t\to 0$, we obtain

$$\|\varphi_T'(\omega) - a(\omega)(\varphi_T(\omega))\| \le \frac{1}{T}d\big(A^{(T)}(\omega)p_0, p_0\big).$$

In particular,

$$\operatorname{speed}(\varphi_T) \le \sup_{\omega\in\Omega}\frac{1}{T}d\big(A^{(T)}(\omega)p_0, p_0\big).$$

The theorem follows by taking $\varphi = \varphi_N$ with sufficiently large N. $\square$

3.6 Creating invariant sections: proof of Theorem D

Proof of Theorem D. By Theorem A, there exists a sequence of sections φ_T such that $\lim_{T\to\infty}\operatorname{speed}(\varphi_T) = 0$. Let $K(w) = K_w$ be the map given by Lemma 3.2. Define

$$\tilde{a}_T(\omega) = a(\omega) + K\big(\varphi_T'(\omega) - a(\omega)(\varphi_T(\omega))\big). \tag{40}$$

(Compare with (35).) Then:

- $\tilde{a}_T(\omega)(\varphi_T(\omega)) = \varphi'_T(\omega)$, that is, φ_T is an invariant section for the cocycle generated by $\tilde{a}_T$.
- for each bounded subset B of H, the sequence $\tilde{a}_T(\omega)(p)$ converges to $a(\omega)(p)$ uniformly with respect to $(\omega, p) \in B \times \Omega$.

Thus the theorem is proved. □

4 Fiber bundle versions

In this section, we consider fiber bundles more general than $\Omega \times H$, and we discuss how to extend some of our results to this setting. We will only deal with the discrete-time case, which is needed for the proof of Theorem 1.5. The continuous-time situation should be similar, but we do not dwell upon it here.

As before, let Ω be a compact Hausdorff space, and let H be a Busemann space. (More hypotheses on H will be imposed along the way.) Endow the group $\mathrm{Isom}(H)$ with the bounded-open topology.

Let Σ be a fiber bundle with base space Ω, fiber H, structural group $\mathrm{Isom}(H)$, and bundle projection $\pi\colon \Sigma \to \Omega$. Thus each fiber $\pi^{-1}(\omega)$ is a metric space (non-canonically) isometric to H. For simplicity, we say that Σ is a *fiber bundle with H fibers.*

Lemma 4.1. *The bundle Σ admits sections, i.e., continuous maps $\varphi\colon \Omega \to \Sigma$ such that $\pi \circ \varphi = \mathrm{id}$.*

Proof. Take a (continuous) partition of unity $\{f_1, \dots, f_n\}$ in Ω such that each f_i is supported in an open set U_i, where $\psi_i\colon U_i \times H \to \pi^{-1}(U_i)$ is a bundle chart. Choose any $p_0 \in H$, and define a map $\varphi\colon \Omega \to \Sigma$ by the formula

$$\varphi(\omega) := \mathrm{bar}_{\pi^{-1}(\omega)} \left(\sum_{i=1}^{n} f_i(\omega) \delta_{\psi_i(\omega, p_0)} \right),$$

where $\mathrm{bar}_{\pi^{-1}(\omega)}$ is the barycenter on the fiber over ω. It follows from the continuity properties of the barycenter (Theorem 2.2) that φ is continuous. □

Let $F\colon \Omega \to \Omega$ be a continuous map. Let $G\colon \Sigma \to \Sigma$ be a bundle map over F, that is, a continuous map G acting isometrically on the fibers and satisfying $\pi \circ G = F \circ \pi$.

Example 4.2. If $A : \Omega \to \mathrm{Isom}(H)$ is a cocycle of isometries, then we can consider the map $G(\omega, v) = (F(\omega), A(\omega)v)$ on the trivial bundle $\Sigma = \Omega \times H$.

In order to keep our formulas similar to the trivial bundle case, we introduce the following notation: For $\omega \in \Omega$ and $n \in \mathbb{Z}_+$, let $A^n(\omega) := G^n(\omega)|\pi^{-1}(\omega)$. This is an isometry from $\pi^{-1}(\omega)$ to $\pi^{-1}(F^n\omega)$. Also denote $A^1(\omega) := A(\omega)$.

The *maximal drift* of G is defined as

$$\mathrm{drift}(G) := \lim_{n\to\infty} \frac{1}{n} \sup_{\omega\in\Omega} d\big(A^{(n)}(\omega)\varphi(\omega), \varphi(F^n\omega)\big), \tag{41}$$

where φ is any (continuous) section of the fiber bundle (whose existence is assured by Lemma 4.1).

The displacement of a section $\varphi : \Omega \to \Sigma$ is defined by the same formula (11).

Proposition 4.3. *For any section φ, we have* $\operatorname{displ}(\varphi) \geq \operatorname{drift}(G)$.

Proof. We have

$$\begin{aligned} d\big(A^{(n)}(\omega)\varphi(\omega), \varphi(F^n\omega)\big) &\leq \sum_{i=0}^{n-1} d\big(A^{(i+1)}(F^{n-i-1}\omega)\varphi(F^{n-i-1}\omega)\varphi(F^{n-i}\omega), A^{(i)}(F^{n-i}\omega)\big) \\ &\leq \sum_{i=0}^{n-1} d\big(A(F^{n-i-1}\omega)\varphi(F^{n-i-1}\omega)\varphi(F^{n-i}\omega)\big) \\ &\leq n \operatorname{displ}(\varphi). \end{aligned}$$

□

Even though it is obvious how to restate Theorems A and B to the fiber bundle setting, we will do it for the reader's convenience:

Theorem E (Fibered version of Theorem A)**.** *Assume that H is a Busemann space. Let Σ be a fiber bundle with H fibers, and let $G\colon \Sigma \to \Sigma$ be a bundle map over F. Then for each $\varepsilon > 0$ there exists a continuous section $\varphi\colon \Omega \to H$ such that* $\operatorname{displ}(\varphi) \leq \operatorname{drift}(G) + \varepsilon$.

Let us say that a subset $B \subset \Sigma$ is *bounded* if for some (and hence any) section $\varphi_0 : \Omega \to \Sigma$, the function $y \in B \mapsto d(y, \varphi_0(\pi(y)))$ is bounded.

Theorem F (Fibered version of Theorem B)**.** *Assume that H is*

- *either a proper Busemann space;*
- *or an Alexandrov space of bounded nonpositive curvature.*

Also assume that H is symmetric. Let Σ be a fiber bundle with H fibers, and let $G\colon \Sigma \to \Sigma$ be a bundle map over F. If $\operatorname{drift}(G) = 0$*, then there is a sequence of bundle maps G_N over F such that:*

- *each G_N has a (continuous) invariant section;*
- *G_N converges to G uniformly on bounded subsets of Σ.*

Let us explain how to adapt the proofs:

Proof of Theorem E. Let $\varphi_1\colon \Omega \to \Sigma$ be a section (whose existence is assured by Lemma 4.1). Modify (30) to:

$$\varphi_N(\omega) := \operatorname{bar}\left(\varphi_1(\omega), [A(\omega)]^{-1}(\varphi_1(F\omega)), \ldots, [A^{(N-1)}(\omega)]^{-1}(\varphi_1(F^{N-1}\omega))\right) .$$

In a similar way as before, one easily checks that this sequence has the required properties. □

Proof of Theorem F. For p_0, p, $q \in H$, let $J_{p_0}(p,q) = J(p,q)$ be defined by formula (34). Notice that $J_{p_0}(p,q)$ also depends continuously on p_0, and the following equivariance property holds:

$$J_{I(p_0)}(I(p), I(q)) = I \circ J_{p_0}(p,q) \circ I^{-1} \quad \text{for every } I \in \operatorname{Isom}(H).$$

As a result, $J_{p_0}(p,q)$ also makes sense if p_0, p, q belong to a same fiber of Σ.

We take a sequence of almost invariant sections φ_N given by Theorem E, and define bundle maps G_N by a formula analogous to (35):

$$\tilde{A}_N(\omega) = J_{\varphi_1(\omega)}\big(A(\omega)\varphi_N(\omega), \varphi_N(F\omega)\big) \circ A(\omega).$$

In a similar way as before, one easily checks that this sequence has the required properties. ☐

5 Invariant conformal structures

Here we prove Theorem 1.5. Before giving the proof, we need to observe a few geometrical facts.

Recall from § 2.9 that the space of ellipsoids in $\mathbb{R}^d$ identifies with Pos_d. In a similar way, the space of *conformal structures* on $\mathbb{R}^d$ identifies with the space SPos_d of positive matrices of determinant 1. This space can be also endowed with a Riemannian metric of nonpositive curvature, and the group $\mathrm{GL}(d,\mathbb{R})$ acts on it by

$$g\colon x \mapsto g(x) := |\det g|^{-2/d} \cdot g \cdot x \cdot g^T.$$

The distance in SPos_d between two elements x, y there is still given by formula (36); this follows from the fact that SPos_d is a totally geodesic subspace of Pos_d (see [Lan, p. 334]).

Proof of Theorem 1.5. Assume $F\colon \Omega \to \Omega$ is a homeomorphism[14] with a unique invariant probability μ, and $\operatorname{supp} \mu = \Omega$. Take any $A \in C^0(\Omega, \mathrm{GL}(d,\mathbb{R}))$; we will explain how to perturb it so that it has the desired properties. First, by [BV], one can perturb A so that the along μ-almost every orbit, the Oseledets splitting is trivial or dominated. Let

$$\mathbb{R}^d = E_1(\omega) \oplus \cdots \oplus E_k(\omega) \tag{42}$$

be the *finest dominated splitting* of the cocycle, that is, the unique[15] everywhere defined global dominated splitting with a maximal number k of bundles (with $k=1$ if there is no dominated splitting).

We claim that for almost every point, there are exactly k different Lyapunov exponents. Indeed, on the one hand, there are at least k different exponents because there is a dominated splitting with k bundles. On the other hand, if there is a positive measure set of points with more than k different Lyapunov exponents, then one can select a point whose orbit is dense on Ω (because the invariant measure has full support) and such that the Oseledets splitting along this orbit is dominated. Since dominated splittings extend to the closure (see [BDV]), one gets a global dominated splitting with more than k bundles; this is a contradiction.

---

[14] Invertibility of the dynamics is needed for having an Oseledets splitting.

[15] See [BDV] for details on finest dominated splittings.

For each $i = 1, \dots, k$, let A_i be the restriction of A to the bundle E_i; then A_i is continuous. Consider the fiber bundle of conformal structures on E_i, and let B_i be the fiber bundle automorphism induced by A_i. It follows from the previous remarks about the space of conformal structures that B_i has uniform sublinear drift. Using Theorem F k times, we perturb A so that each B_i has a continuous invariant section, i.e., each A_i has an invariant conformal structure. Choose a (continuously varying) Riemannian metric on the vector bundle $\Omega \times \mathbb{R}^d$ with respect to which E_i is orthogonal to E_j if $i \neq j$, and whose restriction to each E_i induces the conformal structure obtained above. So there are continuous functions $\varphi_i \colon \Omega \to \mathbb{R}$ such that

$$\|Av\|_{F(\omega)} = e^{\varphi_i(\omega)} \|v\|_\omega \quad \text{for all } v \in E_i(\omega), \tag{43}$$

Let $\lambda_i \in \mathbb{R}$ be the Lyapunov exponent of $\tilde{A}|E_i$. Then the Birkhoff averages of the functions $-\varphi_i + \lambda_i$ tend to zero. By Theorem 1.2 (and its proof), there are functions $\tilde{\varphi}_i$ (close to φ_i) and ψ_i such that

$$-\tilde{\varphi}_i + \lambda_i = \psi_i \circ T - \psi_i.$$

Let $\tilde{A}$ be the perturbation of $\tilde{A}$ defined by

$$\tilde{A}(v) = e^{\tilde{\varphi}_i(\omega) - \varphi_i(\omega)} A(v)$$

Define a new Riemannian metric $\|\cdot\|^*$ with respect to which the splitting (42) is still orthogonal, and such that

$$v \in E_i(\omega) \;\Rightarrow\; \|v\|^*_\omega := e^{\psi_i(\omega)} \|v\|_\omega. \tag{44}$$

Then for all $v \in E_i(\omega)$ we have

$$\begin{aligned} \|\tilde{A}v\|^*_{T(\omega)} &= e^{\tilde{\varphi}_i(\omega) - \varphi_i(\omega) + \psi_i(T\omega)} \|Av\|_{T(\omega)} \\ &= e^{\tilde{\varphi}_i(\omega) + \psi_i(T\omega)} \|v\|_\omega = e^{\tilde{\varphi}_i(\omega) + \psi_i(T\omega) - \psi_i(\omega)} \|v\|^*_\omega = e^{\lambda_i} \|v\|^*_\omega . \end{aligned}$$

This shows that the cocycle $\tilde{A}$ has the desired properties. □

Remark 5.1. It is clear that there is a fibered version of Theorem 1.5, with a nearly identical proof. ◁

Remark 5.2. The conclusion of Theorem 1.5 is obviously false if F is not uniquely ergodic (because if $A \in \mathcal{D}$ then the Lyapunov exponents of A do not depend on the invariant measure). Nevertheless, under the assumption of minimality, one should still be able to obtain invariant conformal structures.

It is claimed in [AB] that for generic matrix cocycles over a minimal homeomorphism of a finite-dimensional compact space, the Oseledets splitting with respect to any invariant probability coincides almost everywhere with the finest dominated splitting. The paper [AB] provides a proof for the case of $\mathrm{SL}(2,\mathbb{R})$ cocycles. Assuming that claim, one can show the following result:

Let $F : \Omega \to \Omega$ be a minimal homeomorphism of a finite-dimensional compact set Ω. Then for every cocycle A in a dense subset $C^0(\Omega, \mathrm{GL}(d,\mathbb{R}))$ there exist:

- *a Riemannian metric on the vector bundle $\Omega \times \mathbb{R}^d$*
- *and a continuous (F, A)-invariant splitting $\mathbb{R}^d = E_1(\omega) \oplus \cdots \oplus E_k(\omega)$ which is orthogonal with respect to the Riemannian metric*

such that the restriction of A to each subbundle E_i is conformal. Moreover the continuous functions $c_i : \Omega \to \mathbb{R}_+$ defined by (21) satisfy $c_1(\omega) > c_2(\omega) > \cdots > c_k(\omega)$ for all $\omega \in \Omega$.

The proof is similar to that of Theorem 1.5, using the claim from [AB] in the place of the result from [BV], and Theorem 1.7 in the place of Theorem 1.2. ◁

A Appendix: The displacement estimate for non-proper spaces

In this appendix, we prove Lemma 2.6. We will actually obtain an explicit formula for the function f. Some preliminaries are needed.

A.1 Angles and more comparisons

Proposition A.1. *Let H be a space of curvature $\leq \kappa$ (resp. $\geq \kappa$, with $\kappa \leq 0$). Let γ_1, γ_2 be two geodesics such that $\gamma_1(0) = \gamma_2(0) = p_0$. For each $t > 0$, $s > 0$, consider the triangle $\triangle(p_0, \gamma_1(t), \gamma_2(s))$, and let $\triangle(\tilde{p}_0, \tilde{p}_{1,t}, \tilde{p}_{2,s})$ be a SSS-comparison triangle in M_κ. Let $\theta_\kappa(t,s)$ be the angle at the vertex $\tilde{p}_0$. Then the function $\theta_\kappa(t,s)$ is monotonically nondecreasing (resp. nonincreasing) with respect to each variable.*

Proof. See [ABN]. □

In particular, the limit of $\theta_\kappa(t,s)$ as $t, s \to 0$ exists. (Notice that the limit actually does not depend on κ.) It is called the *angle* between γ_1 and γ_2 at p_0.

An immediate consequence of Proposition A.1 is that if H has curvature $\leq \kappa$ (resp. $\geq \kappa$) then the angles of a SSS-comparison triangle in M_κ are smaller (resp. greater) than or equal to the corresponding angles for the triangle in H.

If $\triangle(p_1, p_2, p_3)$ is a triangle in H and $\triangle(\tilde{p}_1, \tilde{p}_2, \tilde{p}_3)$ is a triangle in M_κ such that the angles at the vertices p_1 and $\tilde{p}_1$ are equal and the corresponding sides at these vertices have equal lengths (i.e., $d_\kappa(\tilde{p}_1, \tilde{p}_j) = d(p_1, p_j)$ for $j = 2$, 3), then we say that $\triangle(\tilde{p}_1, \tilde{p}_2, \tilde{p}_3)$ is a *SAS-comparison triangle*[16] for $\triangle(p_1, p_2, p_3)$.

---

[16] SAS stands for side-angle-side.

Lemma A.2. *Let H be a uniquely geodesic space of curvature $\geq \kappa$ (where $\kappa \leq 0$). Let $\triangle(p_1, p_2, p_3)$ be a triangle in H, and let $\triangle(\tilde{p}_1, \tilde{p}_2, \tilde{p}_3)$ be a SAS-comparison triangle in M_κ (with equal angles at vertices p_1 and $\tilde{p}_1$). Then:*

1. *The side p_2p_3 is shorter than or has the same length as the side $\tilde{p}_2\tilde{p}_3$.*
2. *If the angles at p_2 and p_3 are less than $\pi/2$, then the angle at p_2 (resp. p_3) is larger than or equal to the angle at $\tilde{p}_2$ (resp. $\tilde{p}_3$).*

Proof. Consider a triangle $\triangle(p_1, p_2, p_3)$ in H (which we assume has curvature $\geq \kappa$). In M_κ, we take a SAS-comparison triangle $\triangle(\tilde{p}_1, \tilde{p}_2, \tilde{p}_3)$ (so that the angles at p_1 and $\tilde{p}_1$ are equal) and a SSS-comparison triangle $\triangle(q_1, q_2, q_3)$. Then the angle at q_i is less than the angle at p_i. Thus, to complete the proof of the lemma, we need the following facts about plane hyperbolic geometry, whose proof we will leave as an exercise.

Claim. *Suppose $\triangle(q_1, q_2, q_3)$ and $\triangle(\tilde{p}_1, \tilde{p}_2, \tilde{p}_3)$ are triangles in M_κ (where $\kappa \leq 0$) so that the angle at vertex r_1 is bigger than the angle at vertex q_1, and the adjacent sides are equal. Then:*

1. *The side $\tilde{p}_2\tilde{p}_3$ is bigger than the side q_2q_3.*
2. *If the angles at $\tilde{p}_2$ and $\tilde{p}_3$ are both less than $\pi/2$, then they are smaller than the angles at q_2 and q_3, respectively.*

□

A.2 The displacement estimate

The following is a more precise version of Lemma 2.6:

Lemma A.3. *Assume that H is a geodesic symmetric space of bounded non-positive curvature in the sense of Alexandrov. Let $\kappa < 0$ be a lower bound for the curvature. Let J be a transvection that translates a geodesic γ by length b. Given $q \in H$, let $s = d(J(q), q)$, $\ell = d(q, \gamma)$, and $\lambda = \sqrt{-\kappa}$. Then*

$$\cosh \lambda s \leq \cosh^2 \frac{\lambda b}{2} \cosh^2 2\lambda\ell + \sinh^2 \frac{\lambda b}{2} \cosh 2\lambda\ell - \cosh \frac{\lambda b}{2} \sinh^2 2\lambda\ell. \tag{45}$$

Proof. Multiplying the metric by a constant, we can assume that $\kappa = -1$, i.e., $\lambda = 1$.

Let J be a transvection that translates a geodesic γ by length b, and let $q \in H$. Let $s = d(J(q), q)$ and $\ell = d(q, \gamma)$. We can assume that $b > 0$ (because for $b = 0$ formula (45) means $s = 0$) and $\ell > 0$ (because for $\ell = 0$ formula (45) means $s \leq b$).

Let p_0 be the point in γ which is closest to q. Let p_1 be the midpoint of p_0 and $J(p_0)$; then $J = \sigma_{p_1} \circ \sigma_{p_0}$. Consider the triangle with vertices p_0, p_1, and $\sigma_{p_0}(q)$. The angle at vertex p_0 is $\pi/2$; let α and β the angles at vertices p_1 and $\sigma_{p_0}(q)$, respectively. See Fig. 3.



Figure 3: Proof of Lemma A.3

Now consider a SAS-comparison triangle in hyperbolic plane M_{-1}, more precisely a triangle in M_{-1} with two sides $b/2$ and ℓ and angle between them $\pi/2$. Let $\tilde{\alpha}$, $\tilde{\beta}$ be the respective angles, and let $\tilde{d}/2$ be the third side. By Lemma A.2, we have

$$d \leq \tilde{d}, \quad \alpha \geq \tilde{\alpha}, \quad \beta \geq \tilde{\beta}.$$

By the law of cosines in M_{-1}, we have:

$$\cosh \frac{\tilde{d}}{2} = \cosh \frac{b}{2} \cosh \ell \,. \tag{46}$$

By the law of sines in M_{-1}, we have:

$$\sinh \frac{\tilde{d}}{2} = \frac{\sinh \ell}{\sin \tilde{\alpha}} \,. \tag{47}$$

By the law of cosines in H (an inequality which comes automatically from the curvature lower bound), we have:

$$\begin{aligned}
\cosh s &\leq \cosh d \cosh 2\ell - \sinh d \sinh 2\ell \cos \beta && \text{(law of cosines)} \\
&\leq \cosh d \cosh 2\ell - \sinh d \sinh 2\ell \sin \alpha && \text{(since } \alpha + \beta \leq \pi/2\text{)} \\
&\leq \cosh d \cosh 2\ell - \sinh d \sinh 2\ell \sin \tilde{\alpha} && \text{(since } \tilde{\alpha} \leq \alpha \leq \pi/2\text{)} \\
&\leq \cosh \tilde{d} \cosh 2\ell - \sinh \tilde{d} \sinh 2\ell \sin \tilde{\alpha} && \text{(since } \tilde{d} \geq d \geq 2\ell\text{)} \\
&= \left(2\cosh^2 \frac{\tilde{d}}{2} - 1\right) \cosh 2\ell - 2 \sinh \frac{\tilde{d}}{2} \cosh \frac{\tilde{d}}{2} \sinh 2\ell \sin \tilde{\alpha}.
\end{aligned}$$

Substituting (46) and (47) and manipulating, we get (45). □

Remark A.4. Fig. 3 is not necessarily contained in a "two-dimensional" totally geodesic subspace. In that particular case, it is possible to show that the following improved version of (45) holds: $\cosh \lambda s \le \cosh \lambda b \cosh^2 \lambda\ell - \sinh^2 \lambda\ell$. Moreover, if $H = M_{-\lambda^2}$, then this becomes an equality, expressing the summit s of a Saccheri quadrilateral as a function of the legs ℓ and the base b; see [BK, p. 104]. ◁

B Appendix: Some lemmas on Killing fields

Proof of part (II) of Lemma 3.1. Let κ be a lower bound for the sectional curvature of H. (Notice that $\kappa > -\infty$ even if H is infinite-dimensional. Indeed, since isometries act transitively on H, it is sufficient to show that sectional curvature is bounded at each point; but this follows directly from the boundedness of the Riemann tensor.) We will show that (II) holds with

$$f(\ell) := \cosh\left(2\sqrt{-\kappa}\cdot\ell\right).$$

Fix p_0, $p \in H$, $v_0 \in T_{p_0}H$. Assume $v_0 \neq 0$, otherwise there is nothing to prove. Let α be the geodesic passing through p_0 with velocity v_0. Let β be a unit-speed geodesic joining p_0 and p. Let $\eta(t) = \xi_{v_0}(\beta(t))$. Then (see [Lan, Prop. 2.2, Ch XIII]) η is a Jacobi field over the geodesic β.

In view of the Rauch–Berger comparison theorem (see [Bi, CE]), in order to show that

$$\|\eta(t)\| \le f(d(\beta(t),\alpha))\|v_0\| \le f(t)\|v_0\|, \tag{48}$$

we need only to consider that case that H is the hyperbolic plane of constant curvature κ. But then it is a simple calculation; actually, in this case, the first inequality in (48) becomes an equality. □

Lemma B.1. *Let H be a symmetric Cartan–Hadamard manifold. For any p_0, $p \in H$, $v \in T_pH$, there exists $\chi \in \mathfrak{m}_{p_0}$ such that $\chi(p) = v$.*

Proof. Let $\ell = d(p, p_0)$. Assume $\ell > 0$, otherwise the claim is trivial. Let $\beta\colon \mathbb{R} \to H$ be the geodesic such that $\beta(0) = p_0$ and $\beta(\ell) = p$. Let $q = \beta(-\ell)$. There is a Jacobi vector field η over β such that

$$\eta(\ell) = v, \quad \eta(-\ell) = 0.$$

(The existence of η follows from the Cartan–Hadamard–McAlpin theorem; see [Lan, § IX.3] and [Lan, Thrm. IX.3.1].) We claim that $\xi_{2\eta(0)}$ is the sought-after Killing field.

Let $\sigma = \sigma_{p_0}$ be the symmetry at p_0; then $\sigma(\beta(t)) = \beta(-t)$. Let ζ be the Jacobi field over β obtained by pushing-forward η by σ, that is,

$$\zeta(t) = T\sigma(\beta(-t)) \cdot \eta(-t).$$



Figure 4: The Jacobi field η, its reflection ζ, and the field $\chi = \eta - \zeta$.

Consider the Jacobi field $\chi = \eta - \zeta$. (See Fig. 4.)

The isometry σ preserves covariant derivative, so

$$\nabla_{\beta'}\zeta(t) = -T\sigma(\beta(-t)) \cdot \nabla_{\beta'}\eta(-t).$$

Taking $t = 0$, we get $\nabla_{\beta'}\chi(0) = 0$. As a consequence (use [Lan, Prop. XIII.5.6]), the Jacobi field χ over β can be extended to a Killing field $\chi \in \mathfrak{m}_{p_0}$. This is exactly $\xi_{\chi(0)} = \xi_{2\eta(0)}$. □

C Appendix: Differentiability of the Cartan barycenter

In this appendix, we give the proof of Lemma 3.4.

Assume H is a Cartan–Hadamard manifold. Let μ be a probability measure of bounded support (p.o.b.s.) on H, and let $f = f_\mu$ be given by (29).

Lemma C.1. *The gradient vector field*[17] *of f is given by:*

$$\operatorname{grad} f(p) = -\int_H \exp_p^{-1}(q)\, \mathrm{d}\mu(q)\,.$$

Proof. See [BK, p. 132]. □

Lemma C.2. *For each p, the linear map*

$$L\colon v \in T_pM \mapsto \nabla_v \operatorname{grad} f(p) \in T_pM$$

(given by covariant derivative) is bounded and has a bounded inverse.

---

[17] Defined by the relation $Tf_p(v) = \langle \operatorname{grad} f(p), v\rangle$.

Proof. We have $\operatorname{grad} f = \int_H \xi_q \, \mathrm{d}\mu(q)$, where ξ_q is the vector field $\xi_q(p) = -\exp_p^{-1}(q)$. For each q and p, the linear map $v \in T_pM \mapsto \nabla_v \xi_q(p) \in T_pM$ is symmetric and $\geq \mathrm{id}$; see [Karc, p. 172, 188]. Integrating over q, we conclude that L is self-adjoint and $\geq \mathrm{id}$. Now we need the following fact:

Claim. *Let L be a bounded symmetric operator on real Hilbert space such that $L \geq \mathrm{id}$. Then L is invertible, with a bounded inverse.*

Proof of the Claim. Let $c > 0$ and estimate

$$\|(\mathrm{id} - cL)v\|^2 = \langle v, v\rangle - 2c\langle Lv, v\rangle + c^2\langle Lv, Lv\rangle \leq \left(1 - 2c + c^2\|L\|^2\right)\|v\|^2 .$$

So if c is small enough, then $\|\mathrm{id} - cL\| < 1$. In particular, $-cL$ is invertible, and so is L. ☐

The lemma follows. ☐

Proof of Lemma 3.4. Define a one-parameter family of vector fields ξ_t $(t \in I)$ on H by

$$\xi_t(p) = -\frac{1}{T}\int_0^T \exp_p^{-1}(h(t,s))\,\mathrm{d}s\,.$$

Then $\bar{h}(t)$ is the unique solution of $\xi_t(\bar{h}(t)) = 0$. To deduce differentiability of $\bar{h}$ from the Implicit Function Theorem, it is sufficient to check that for each $t \in H$ and $p \in H$, the linear map $v \in T_pM \mapsto \nabla_v \xi_t(p) \in T_pM$ (given by covariant derivative) is invertible (with a bounded inverse) – see [BK, p. 143] for details. But this was proved in Lemma C.2 above. ☐

Remark C.3. Another approach to the differentiability of the barycenter is to consider barycenters in the tangent bundle TH; see [AL]. ◁

References

[ABN] A.D. Aleksandrov, V.N. Berestovskiĭ & I.G. Nikolaev. Generalized Riemannian spaces. *Russian Math. Surveys* **41**, no. **3** (1986), 1–54. MR0854238 (Cited on page 29.)

[AL] M. Arnaudon & X.-M. Li. Barycenters of measures transported by stochastic flows. *Ann. Probab.* **33**, no. **4** (2005), 1509–1543. MR2150197 (Cited on pages 14 and 34.)

[ANT] T. Austin, A. Naor & R. Tessera. Sharp quantitative nonembeddability of the Heisenberg group into superreflexive Banach spaces. Preprint arXiv:1007.4238 (Cited on page 12.)

[AB] A. Avila & J. Bochi. A uniform dichotomy for generic cocycles over a minimal base. *Bull. Soc. Math. France* **135**, no. **3** (2007), 407–417. MR2430187 (Cited on pages 7, 28, and 29.)

[ABD1] A. Avila, J. Bochi & D. Damanik. Cantor spectrum for Schrödinger operators with potentials arising from generalized skew-shifts. *Duke Math. J.* **146**, no. **2** (2009), 253–280. MR2477761 (Cited on pages 4, 18, and 23.)

[ABD2] ______. Opening gaps in the spectrum of strictly ergodic Schrdinger operators. *J. Eur. Math. Soc.* **14**, no. **1** (2012), 61–106. (Cited on pages 4, 11, 14, and 19.)

[AK] A. Avila, A. Kocsard. Cohomological equations and invariant distributions for minimal circle diffeomorphisms. *Duke Math. J.* **158**, no. **2** (2011), 501–536. MR2805066 (Cited on page 11.)

[Bi] L. Biliotti. The exponential map of a weak Riemannian Hilbert manifold. *Illinois J. Math.* **48**, no. **4** (2004), 1191–1206. MR2113672 (Cited on page 32.)

[BV] J. Bochi & M. Viana. The Lyapunov exponents of generic volume preserving and symplectic maps. *Ann. Math.* **161**, no. **3** (2005), 1423–1485. MR2180404 (Cited on pages 4, 10, 27, and 29.)

[BDV] C. Bonatti, L. J. Díaz & M. Viana. *Dynamics beyond uniform hyperbolicity.* Springer (2005). MR2105774. (Cited on pages 27 and 28.)

[BH] M.R. Bridson & A. Haefliger. *Metric spaces of non-positive curvature.* Grundlehren der mathematischen Wissenschaften **319**, Springer (1999). MR1744486 (Cited on pages 17 and 20.)

[BK] P. Buser & H. Karcher. Gromov's almost flat manifolds. *Astérisque* **81**, Soc. Math. France (1981). MR0619537 (Cited on pages 14, 32, 33, and 34.)

[Ca] E. Cartan. *Leçons sur la géométrie des espaces de Riemann.* Gauthier-Villars (1928). (Cited on page 14.)

[CE] J. Cheeger & D.G. Ebin. *Comparison theorems in Riemannian geometry.* Revised reprint of the 1975 original. AMS Chelsea Publishing (2008). MR2394158 (Cited on page 32.)

[CLT] G. Contreras, A.O. Lopes, P. Thieullen. Lyapunov minimizing measures for expanding maps of the circle. *Erg. Theory Dyn. Sys.* **21**, no. **5**, 1379–1409. MR1855838 (Cited on page 11.)

[CPR] G. Corach, H. Porta & L. Recht. A geometric interpretation of Segal's inequality $\|e^{X+Y}\| \leq \|e^{X/2}e^{Y}e^{X/2}\|$. *Proc. Amer. Math. Soc.* **115**, no. **1**, (1992) 229–231. MR1075945 (Cited on page 20.)

[CNP1] D. Coronel, A. Navas & M. Ponce. Bounded orbits versus invariant curves for cocycles of affine isometries over a minimal dynamics. Preprint arXiv:1101.3523 (Cited on page 3.)

[CNP2] ______. On the dynamics of non-reducible cylindrical vortices. To appear in *J. London Math. Soc.* Preprint arXiv:1101.3526 (Cited on page 5.)

[EH] A. Es-Sahib & H. Heinich. Barycentre canonique pour un espace métrique à courbure négative. Séminaire de Probabilités XXXIII. *Lecture Notes in Math.* **1709**, Springer, Berlin (1999), 355–370. MR1768010 (Cited on page 14.)

[Fu] A. Furman. On the multiplicative ergodic theorem for uniquely ergodic systems. *Ann. Inst. H. Poincaré Probab. Statist.* **33** (1997), 797–815. MR1484541 (Cited on page 7.)

[GL] E. Garibaldi & A.O. Lopes. On the Aubry-Mather theory for symbolic dynamics. *Erg. Theory Dyn. Sys.* **28**, no. **3**, 791–815. MR2422016 (Cited on page 11.)

[He] S. Helgason. *Differential Geometry, Lie Groups, and Symmetric Spaces.* Corrected reprint of the 1978 original. Graduate Studies in Mathematics **34**, Amer. Math. Soc. (2001). MR1834454 (Cited on page 15.)

[Je] O. Jenkinson. Ergodic optimization. *Discrete Contin. Dyn. Sys.* **15**, no. **1** (2006), 197–224. MR2191393 (Cited on pages 10 and 11.)

[Jo] J. Jost. *Nonpositive curvature: geometric and analytic aspects.* Lectures in Mathematics – ETH Zürich, Birkhäuser (1997). MR1451625 (Cited on page 14.)

[Kai] V.A. Kaimanovich. Lyapunov exponents, symmetric spaces and a multiplicative ergodic theorem for semisimple Lie groups. *J. Soviet Math.* **47**, no. **2** (1989), 2387–2398. MR0947327 (Cited on page 6.)

[KalS] B. Kalinin & V. Sadovskaya. Linear cocycles over hyperbolic systems and criteria of conformality. *J. Modern Dyn.* **4**, no. **3** (2010), 419–441. MR2729329 (Cited on page 10.)

[Karc] H. Karcher. Riemannian comparison constructions. *Global differential geometry*, 170–222, MAA Stud. Math. **27**, Math. Assoc. America, 1989. MR1013810 (Cited on page 34.)

[KarL] A. Karlsson, F. Ledrappier. Noncommutative ergodic theorems. *Geometry, rigidity, and group actions* (Farb and Fisher, eds.). Chicago Lectures in Mathematics, Chicago Univ. Press (2011), 396–418. MR2807838 (Cited on page 6.)

[KarM] A. Karlsson, G.A. Margulis. A multiplicative ergodic theorem and nonpositively curved spaces. *Comm. Math. Phys.* **208** (1999), 107–123. MR1729880 (Cited on page 6.)

[KatH] A. Katok & B. Hasselbatt. *Introduction to the modern theory of dynamical systems.* Cambridge Univ. Press, 1996. MR1326374 (Cited on page 3.)

[Kat] A. Katok. Cocycles, cohomology and combinatorial constructions in ergodic theory. In collaboration with E. A. Robinson, Jr. *Smooth ergodic theory and its applications* (Katok, de la Llave, Pesin, and Weiss, eds.) *Proc. Sympos. Pure Math.*, **69**, Amer. Math. Soc. (2001), 107–173. MR1858535 (Cited on pages 5 and 11.)

[Lan] S. Lang. *Fundamentals of Differential Geometry.* Graduate Texts in Mathematics **191**, Springer (1999). MR1666820 (Cited on pages 19, 20, 21, 22, 23, 27, 32, and 33.)

[Lar] G. Larotonda. Nonpositive curvature: a geometrical approach to Hilbert-Schmidt operators. *Differential Geom. Appl.* **25** (2007), 679–700. MR2373944 (Cited on page 20.)

[Mo] I.D. Morris. A rapidly-converging lower bound for the joint spectral radius via multiplicative ergodic theory. *Adv. Math.* **225** (2010), 3425–3445. MR2729011 (Cited on page 3.)

[MOP1] J. Moulin Ollagnier & D. Pinchon. Mesures quasi invariantes à dérivée continue. *C. R. Acad. Sci. Paris Sér. A-B* **282** no. **23** (1976), Aii, A1371–A1373. MR0422579 (Cited on page 12.)

[MOP2] ______. Systèmes dynamiques topologiques I. Étude des limites de cobords. *Bull. Soc. Math. France* **105** (1977), 405–414. MR0498967 (Cited on pages 5 and 11.)

[Na] A. Navas. An L^1 ergodic theorem with values in a nonpositively curved space via a canonical barycenter map. To appear in *Erg. Theory Dyn. Sys.* Preprint arXiv:1104.5188 (Cited on page 14.)

[Pa] A. Papadopoulos. *Metric spaces, convexity and nonpositive curvature.* IRMA Lectures in Mathematics and Theoretical Physics **6**, EMS (2005). MR2132506 (Cited on page 16.)

[Pau] S.D. Pauls. The large scale geometry of nilpotent Lie groups. *Comm. Anal. Geom.* **9**, no. **5** (2001), 951982. MR1883722 (Cited on page 12.)

[Qu] A.N. Quas. Rigidity of continuous coboundaries. *Bull. London Math. Soc.* **29** (1997), 595–600. MR1458721 (Cited on page 3.)

[Schr] S.J. Schreiber. On growth rates of subadditive functions for semiflows. *J. Differential Equations* **148** (1998), 334–350. MR1643183 (Cited on page 7.)

[Schw] S. Schwartzman. Asymptotic cycles. *Ann. Math.* **66** (1957), 270–284. MR0088720 (Cited on page 9.)

[St] K.-T. Sturm. Probability measures on metric spaces of nonpositive curvature. *Heat kernels and analysis on manifolds, graphs, and metric spaces.* (Auscher, Coulhon and Grigor'yan, eds.) Contemporary Math. **338** (2003), 357–390. MR2039961 (Cited on pages 14 and 17.)

[SS] R. Sturman & J. Stark. Semi-uniform ergodic theorems and applications to forced systems. *Nonlinearity* **13** (2000), 113–143. MR1734626 (Cited on page 7.)

[Vi] C. Villani. *Optimal transport, old and new.* Grundlehren der Mathematischen Wissenschaften **338**, Springer (2009). MR2459454 (Cited on page 13.)

Jairo Bochi
Pontifícia Universidade Católica do Rio de Janeiro (PUC–Rio)
Rua Marquês de S. Vicente, 225. Rio de Janeiro, Brazil.
jairo@mat.puc-rio.br
www.mat.puc-rio.br/∼jairo

Andrés Navas
Universidad de Santiago
Alameda 3363, Estación Central. Santiago, Chile.
andres.navas@usach.cl